

João Galamba Correia

From: onbehalfof+dalton+rsc.org@manuscriptcentral.com on behalf of Dalton Transactions <onbehalfof+dalton+rsc.org@manuscriptcentral.com>
Sent: 30 de maio de 2017 11:22
To: João Galamba Correia
Subject: Decision on submission to Dalton Transactions - DT-ART-03-2017-001146.R1

30-May-2017

Dear Dr Correia:

Manuscript ID: DT-ART-03-2017-001146.R1

TITLE: Technetium-99m complexes of L-arginine derivatives for targeting amino acid transporters

Thank you for submitting your revised manuscript to Dalton Transactions. After considering the changes you have made, **I am pleased to accept your manuscript for publication in its current form in the Dalton Transactions themed issue on Frontiers in Radionuclide Imaging and Therapy.** I have copied any final comments from the reviewer(s) below.

Please note that if you have requested Accepted Manuscript publication we will publish your article shortly and send you an email once it is available online. We will also email you information on how to access your Dalton Transactions article proofs shortly.

If you would like to publish your article open access, once your article is accepted for publication, please complete our open access application form and agree to a Creative Commons licence. The form and further information can be found at:

<http://rsc.li/open-access>

Your funder or research institute may also have certain requirements or mandates for open access publication. If you are funded by Wellcome or are from a Max Planck institute, please DO NOT complete the standard licence to publish and see our funder requirements guidance for more information:

<http://www.rsc.org/journals-books-databases/open-access/#requirements>

You can highlight your article and the work of your group on the back cover of Dalton Transactions, if you are interested in this opportunity please contact me for more information.

As an author you are entitled to a 25% discount on books published by the Royal Society of Chemistry. To receive this discount, enter the promotional code JLTH25 when purchasing from our online bookshop (pubs.rsc.org/bookshop). Please contact booksales@rsc.org if you have any problems.

Discover more Royal Society of Chemistry author services and benefits here:

<http://www.rsc.org/journals-books-databases/about-journals/#author-benefits>

Thank you for publishing with Dalton Transactions, a journal published by the Royal Society of Chemistry – the world's leading chemistry community, advancing excellence in the chemical sciences.

With best wishes,

Caroline Knapp

Dr Caroline Knapp
Publishing Editor, Dalton Transactions
Royal Society of Chemistry
Thomas Graham House,
Science Park, Milton Road
Cambridge, CB4 0WF, UK
Tel: +44 (0)1223 432531

Follow the Royal Society of Chemistry:
www.rsc.org/follow

Winner of The Queen's Award for Enterprise, International Trade 2013

REVIEWER REPORT(S):

Referee: 1

Comments to the Author

The authors have addressed all the comments by the reviewers to acceptable and publishable levels.

DISCLAIMER:

This communication (including any attachments) is intended for the use of the addressee only and may contain confidential, privileged or copyright material. It may not be relied upon or disclosed to any other person without the consent of The Royal Society of Chemistry. If you have received it in error, you must not copy or show it to anyone; please contact us immediately by replying to this email and highlighting the error. Any advice given by The Royal Society of Chemistry has been carefully formulated but is necessarily based on the information available, and The Royal Society of Chemistry cannot be held responsible for accuracy or completeness. In this respect, any views or opinions presented in this email are solely those of the author and may not represent those of The Royal Society of Chemistry.

The Royal Society of Chemistry owes no duty of care and shall not be liable for any resulting damage or loss as a result of the use of this email and/or attachments. The Royal Society of Chemistry acknowledges that a disclaimer cannot restrict liability at law for personal injury or death arising through a finding of negligence. The Royal Society of Chemistry does not warrant that its emails or attachments are Virus-free: Please rely on your own screening. The Royal Society of Chemistry is a charity, registered in England and Wales, Number 207890, and a company incorporated in England by Royal Charter (Registered No. RC000524) Registered office: Burlington House, Piccadilly, London W1J 0BA, Telephone: 0207 4378 6556, Facsimile: 0207 4490 3393 (Head Office).

Technetium-99m complexes of L-arginine derivatives for targeting amino acid transporters

Maurício Morais,^{1,2,‡} Vera F. C. Ferreira,^{1,‡} Flávio Figueira,¹ Filipa Mendes,¹ Paula Raposinho,¹
Isabel Santos,¹ Bruno L. Oliveira,^{1,3} João D. G. Correia^{1,*}

¹Centro de Ciências e Tecnologias Nucleares, Instituto Superior Técnico, Universidade de Lisboa, Estrada Nacional 10 (km 139,7), 2695-066 Bobadela LRS, Portugal

²Current address: Division of Imaging Sciences and Biomedical Engineering, King's College London, 4th Floor Lambeth Wing, St Thomas' Hospital, London SE1 7EH, United Kingdom

³Current address: Department of Chemistry, University of Cambridge, Lensfield Road, Cambridge, United Kingdom

Submitted as a full article to: *Dalton Trans.*, *Frontiers in Radionuclide Imaging and Therapy* themed issue

Keywords: Amino acid transporters, Arginine, Cancer, Imaging, Rhenium, Technetium

* Corresponding author: João D. G. Correia, Tel.: +351 21 994 62 33

E-mail address: jgalamba@ctn.tecnico.ulisboa.pt

‡ These authors contributed equally to the article. Author contributions: J.D.G.C and B.L.O. conceived the experimental strategy; F.F. and M.M. performed the synthesis, radiolabelling and enzymatic studies; V.F.C.F. conducted cell culture, internalization studies and western blotting; P.R. and F.M. provided intellectual input on the cellular studies, enzymatic studies in cells and western blotting; I.S., B.L.O and J.D.G.C prepared the draft and final version of the manuscript. B.L.O and J.D.G.C are co-senior authors of this work.

Abstract

Although relevant from the clinical point of view, radiotracers targeting cationic amino acid transporters are relatively unexplored and, in particular, no metal-based radiotracers are known. The rare examples of complexes recognized by amino acid transporters, namely by the Na⁺-independent neutral L-type amino acid transporter 1 (LAT1), are ^{99m}Tc(I)/Re(I) compounds. Herein, we describe conjugates comprising a pyrazolyl-diamine chelating unit and the cationic amino acid L-arginine (L-Arg) linked by a propyl (**L**¹) or hexyl linker (**L**²), which allowed the preparation of stable complexes of the type *fac*-[^{99m}Tc(CO)₃(κ³-L)] (**Tc1**, L = **L**¹; **Tc2**, L = **L**²) and of the respective surrogates **Re1** and **Re2**. Interestingly, complex **Tc2** exhibited moderate levels of time-dependent internalization in three human tumoural cell lines, with approximately 3 % of total applied activity internalized, corresponding to 21 % of cell-associated activity. A putative mechanism of retention in the cytoplasm of cells could be the interaction of the complex with inducible nitric oxide synthase (iNOS), which is the enzyme responsible for the catalytic oxidation of L-Arg to citrulline and nitric oxide. However, the surrogate complex **Re2** does not recognize iNOS, as demonstrated by the *in vitro* assays with purified iNOS and in studies with lipopolysaccharide(LPS)-activated macrophages. Preliminary mechanistic studies suggest that the internalization of **Tc2** is linked to the cationic amino acid transporters, namely system y⁺. This finding might open the way towards the development of novel families of metal-based radiotracers for probing metabolically active cancer cells.

Introduction

The majority of the Positron Emission Tomography (PET) imaging procedures rely on the use of the glucose analogue 2-[¹⁸F]fluoro-2-deoxy-D-glucose ([¹⁸F]FDG), which enters the cells via membrane glucose transporters, where it undergoes phosphorylation and is irreversibly trapped.^{1, 2} Accumulation of this radiopharmaceutical in tumoural cells is mainly due to the upregulation of glucose transport and glycolysis. However, [¹⁸F]FDG presents some limitations, including limited visualization of brain tumors, low or variable uptake in some tumors types (e.g. prostate cancer and neuroendocrine tumors) and increased accumulation in inflammatory lesions.^{3, 4} Thus, radiolabeled amino acids, which target increased rates of amino acid transport in cancer cells, have been considered as alternatives to overcome some of those limitations.^{5, 6} Indeed, they are accepted as tracers for imaging the upregulated metabolism linked to several hallmarks of cancer.

The transport of amino acids across the plasma membrane into the cytoplasm of mammalian cells is mediated by membrane-bound transport systems that present varying substrate specificities, pH and sodium dependence, and regulatory mechanisms.⁶⁻⁸ Most of the work performed so far was based on radiolabeled amino acids for targeting the “L amino acid transport system”, which preferentially transports amino acids with neutral side chains like L-Leu, L-Tyr, and L-Phe.^{9, 10} Relevant tracers from this class include L-¹¹C-methionine and O-(2-¹⁸F-fluorethyl)-L-Tyrosine (¹⁸F-FET) for imaging brain tumors, and 6-¹⁸F-fluoro-3,4-dihydroxy-L-phenylalanine (¹⁸F-FDOPA) for imaging of neuroendocrine tumors. In the past few years there has been a growing interest in the design of tracers for targeting other amino acid transporters, including “system A”, glutamine, glutamate and cationic amino acid transporters.⁹ The transport of basic amino acids (L-Lys, L-Arg and L-His) is mediated by sodium-independent and sodium-dependent transporter systems, which include the cationic amino acid transporter (CAT, system y⁺) family, system y⁺L, b^{0,+}AT and ATB^{0,+}.

The amino acid L-Arg is the precursor for relevant metabolic pathways such as agmatine, creatine, urea, and nitric oxide (NO) synthesis.¹¹ The latter is a mammalian signaling molecule biosynthesized by NO synthases (NOS) with high relevance in physiological (e.g. neuronal transmission) and pathological processes (e.g. cancer and neurological disorders).¹²⁻¹⁴ Additionally, L-Arg is a relevant signaling molecule that regulates essential cellular functions such as protein synthesis, apoptosis and growth.¹¹ It also plays an important role in cells lacking argininosuccinate synthase 1 (ASS1), one of the key urea cycle enzymes that is absent in many tumors, suggesting that tumoral ASS1 deficiency may be both a prognostic biomarker and predictor of sensitivity to arginine deprivation therapy.¹⁵ Brought together the pathophysiological roles of L-Arg and the fact that certain cancer cells overexpress cationic amino acid transporters such as ATB^{0,+} or CAT1,¹⁶⁻²⁰ indicate that radiolabelled L-Arg derivatives hold potential for cancer imaging. Moreover, radiolabelled L-Arg derivatives could be envisaged as imaging biomarkers for predicting and monitoring response to arginine deprivation therapy.


Although potentially relevant from the clinical point of view, radiotracers targeting cationic amino acid transporters are relatively unexplored and, in particular, no metal-based radiotracers are known.^{6,9} So far, the only examples of metal complexes recognized by amino acid transporters and actively internalized into cancer cells, more specifically through the L-type LAT1, are ^{99m}Tc(I)/Re(I) complexes.²¹ One of the main advantages of using ^{99m}Tc-based complexes, compared to using cyclotron-produced radionuclides such as ¹¹C or ¹⁸F, relies on fact that ^{99m}Tc is easily available in nuclear medicine centres worldwide through ⁹⁹Mo/^{99m}Tc generators. Moreover, it is considered to burden a low dose of radiation to patients. Considering our previous work in the design and biological evaluation of novel Tc(I)/Re(I) complexes with pendant L-Arg derivatives for visualization of NO/NOS-related tumors by SPECT imaging,²²⁻²⁶ we describe herein the (radio)synthesis, biological evaluation and

preliminary mechanistic studies of novel $^{99m}\text{Tc(I)}$ -labelled L-Arg derivatives useful for imaging metabolically active cells.

Results and discussion

Complexes of the type $\text{fac-}[\text{M}(\text{CO})_3(\text{k}^3\text{-L})]$ ($\text{M} = \text{Re}/^{99m}\text{Tc}$, $\text{L} = \text{L}^1$ and L^2)

We have prepared and fully characterized conjugates L^1 and L^2 containing L-Arg following previously described procedures.²² Besides the pendant amino acid moiety, the conjugates present a pyrazoly-diamine chelating unit known to stabilize efficiently the organometallic core $\text{fac-}[\text{M}(\text{CO})_3]^+$ ($\text{M} = ^{99m}\text{Tc}$ or Re), which already allowed the radiolabeling of various molecules with biological relevance.^{23, 27} In brief, the conjugates were prepared upon conjugation of the Boc-protected precursors **Pz-C₃NH₂(Boc)** and **Pz-C₆NH₂(Boc)** to *N*- α -Boc-L-Arg (Scheme 1), respectively, using standard coupling reagents and conditions (HBTU, Et₃N, 2 h, room temperature), followed by hydrolysis of the protecting groups with trifluoroacetic acid (TFA).



Scheme 1. Synthesis of L^1 , L^2 , Tc1/Re1 and Tc2/Re2 (Identification system for NMR assignments is displayed for the rhenium complexes). i) Et_3N , N - α -Boc-L-Arg, HBTU, 2 h, r.t.; ii) CH_2Cl_2 -TFA, 3 h, r.t.; iii) $[\text{M}(\text{CO})_3(\text{H}_2\text{O})_3]^+$ ($M = \text{Re}$, ${}^{99\text{m}}\text{Tc}$), H_2O , 100°C .

Conjugates L^1 and L^2 were obtained in high purity ($\geq 98\%$ ascertained by RP-HPLC, Figure S1) as stable colorless oils after purification by semi-preparative reversed phase high performance liquid chromatography (RP-HPLC). The conjugates were fully characterized by ${}^1\text{H}/{}^{13}\text{C}$ NMR (Figure S2 and Figure S3) and IR spectroscopy as well as electrospray ionization mass spectrometry.

The radioactive complexes $fac-[^{99m}\text{Tc}(\text{CO})_3(\text{k}^3\text{-L})]$ (**Tc1**, L = **L¹**; **Tc2**, L = **L²**, Scheme 1) were prepared in high radiochemical yield and radiochemical purity (> 95%) upon reaction of **L¹** or **L²** with the precursor $fac-[^{99m}\text{Tc}(\text{CO})_3(\text{H}_2\text{O})_3]^+$ (Figure S4). The latter was prepared by addition of $\text{Na}[^{99m}\text{TcO}_4]$, eluted from a $^{99}\text{Mo}/^{99m}\text{Tc}$ generator with saline solution, to an IsoLink kit (Mallinckrodt, Covidien) available for research purposes, and heating (95 °C) for 20 min. The high stability of the resulting final complexes was demonstrated by incubation with a 100-fold excess of coordinating amino acids such as histidine (Figure S5) or cysteine. No relevant degradation or transchelation were detected by RP-HPLC after incubation at 37°C for up to 6 h in line with earlier results obtained for complexes stabilized by the same chelating unit.^{22, 27, 28} Additional stability studies demonstrated that the complexes are also stable in human plasma. The dilute nature of solutions of ^{99m}Tc complexes (ca. 10^{-9} - 10^{-12} M) hampers their structural characterization by the usual analytical methods in chemistry. The more straightforward way to overcome this limitation is to compare the chromatographic behavior of ^{99m}Tc complexes with that of the surrogate rhenium complexes prepared at the “macroscopic” scale, since technetium and rhenium, transition metals of group 7 of the periodic table, share similar coordination chemistry. Thus, the chemical identity of **Tc1** and **Tc2** was confirmed by comparing their RP-HPLC radioactive traces (γ detection) with the UV-Vis traces of the surrogate complexes **Re1** and **Re2** (Figure S6). These complexes were synthesized upon reaction of **L¹** and **L²** with $fac\text{-Re}(\text{CO})_3(\text{H}_2\text{O})_3]^+$ in refluxing water (Scheme 1), and were obtained in moderate yields (35 – 75%) after purification by semi-preparative RP-HPLC. **Re1** and **Re2** were fully characterized by ESI-MS, IR and $^1\text{H}/^{13}\text{C}$ NMR spectroscopy. The data collected support the proposed structure and the tridentate coordination mode of the pyrazolyl-diamine chelating unit, comparing well with similar complexes previously described by our group^{22, 27, 29}.

Cellular uptake studies

Aimed at predicting the *in vivo* tumor-targeting properties of **Tc1** and **Tc2** and to assess their ability to be recognized and internalized as specific substrates by the transporters of L-Arg, we have performed uptake studies in a panel of human tumoral cell lines, more specifically HeLa cervical cancer, A375 melanoma, MDA-MB-231 breast cancer and PC3 prostate cancer cell lines. The results of the cellular uptake as a function of the incubation time are presented in Figure 1.





Figure 1. Cellular uptake of **Tc1** and **Tc2** in human cancer cell lines at 37°C.

Complex **Tc2** displayed a remarkably higher uptake than **Tc1**, with values between 11.4 ± 0.7 % (HeLa cell line) and 15.1 ± 0.3 % (A375 cell line) after 3 h incubation. We speculate that the difference observed between the uptake values for the two radioconjugates might be assigned to the presence of a longer alkyl chain in **Tc2** (hexyl linker) than in **Tc1** (propyl linker), assuming that these arginine derivatives are substrates of the Na^+ -independent transport system γ^+ , which has been postulated to be the major entry route for cationic amino acids, L-Arg included, in most cells.

Indeed, it has been reported that for system γ^+ , CAT proteins have a higher affinity for cationic amino acids with a longer carbon backbone: homoarginine > arginine > lysine >

ornithine > 2,4-diamino-*n*-butyric acid.⁸ Other clue suggesting the involvement of the system γ^+ in the cellular uptake of **Tc1** and **Tc2** is the fact that analogue radioconjugates with pendant N^{ω} -NO₂-L-arginine moieties (**Tc3** - **Tc5**, Scheme 2), previously prepared for targeting NOS,^{22, 28} are not taken up by the cell lines tested (Figure 2).



Scheme 2. Molecular structures of **Tc3** - **Tc5**.

In fact, it has been described that inhibitors of NOS such as N^{ω} -NO₂-L-arginine methyl ester or *N*-methyl arginine competitively inhibit arginine transport across cell membranes due to interaction with system γ^+ .³⁰





Figure 2. Cellular uptake of the radioconjugates **Tc3** (A), **Tc4** (B) and **Tc5** (C) in various human cancer cell lines at 37°C.

The remarkable cellular uptake results obtained with **Tc2** prompted us to deepen these studies in an attempt to unveil the most likely cellular uptake mechanism, using both the radioactive complex **Tc2** and the “cold” surrogate **Re2**. Therefore, internalization studies in HeLa, A375 and MDA-MB-231 cancer cell lines have been performed, and the results are displayed in Figure 3A.






Figure 3. A) Internalized and surface-bound **Tc2** in HeLa, A375 and MDA-MB-231 cancer cell lines at 37°C, expressed as a percentage of total applied activity. **B)** Internalized and surface-bound **Tc2** after 2 h in the presence of lysine at 37°C, expressed as a percentage of cellular uptake in the absence of L-Lys.

Tc2 exhibits moderate levels of time-dependent internalization with the highest values being reached at 4 h in all cell lines: 2.7 ± 0.2 % (A375), 3.0 ± 0.3 % (MDA-MB-231) and 3.2 ± 0.3 % (HeLa) of the total applied activity internalized, corresponding to 21.3 ± 0.8 %, 21.6 ± 1.3 % and 22.5 ± 1.0 % of the cell-associated activity, respectively.

Aimed at clarifying the role of the transporters of cationic amino acids in the uptake and internalization of **Tc2**, and considering that L-Lys competes with L-Arg for common transport systems,^{11, 31, 32} we have performed internalization studies of **Tc2** with co-incubation with L-Lys in the same cell lines (Figure 3B). Although moderate, the inhibition of cell surface-bound **Tc2** (17%, 25% and 18% for HeLa, MDA-MB-231 and A375 cell lines, respectively) and, more importantly, the inhibition of internalized **Tc2** (20%, 29% and 36%, respectively) suggest involvement of the transporters of cationic amino acids in the cell uptake mechanism of **Tc2**.

Furthermore, we have also studied the contribution of the CAT family (system γ^+), more specifically of CAT1, upregulated in various cancers,^{19, 20} in the specific uptake of **Tc2**. Firstly,

we have confirmed the expression of CAT1 in the selected cell lines by Western blot analysis of protein extracts using an anti-CAT1 antibody (Figure 4).




Figure 4. Evaluation of CAT1 expression in HeLa cervical cancer, A375 melanoma, MDA-MB-231 breast cancer and PC3 prostate cancer cell lines. Bands corresponding to unglycosylated (ca. 70 kDa) form and glycosylated form (ca. 90 kDa) of CAT1 are observed. Actin was used as an internal loading control.

Then, we performed cell uptake assays in the presence of the cysteine-modifying agent *N*-ethylmaleimide (NEM), which has been recently shown to inhibit selectively all human CATs (system γ^+), but no other transporters of cationic amino acid transporters such as γ^+ LAT isoform.³³ The results of the cell uptake studies are presented in Figure 5.




Figure 5. A) Cellular uptake of **Tc2** in the A375 melanoma cell line at different time points at 37°C in the presence of N-ethylmaleimide (NEM, 5 mM). **B)** Inhibition of cellular uptake by NEM expressed in percentage.

Under the conditions tested, the cellular uptake of **Tc2** is inhibited by NEM in 10.5 %-20.9 % after 5 to 60 minutes incubation (Figure 5B), which supports our hypothesis of CAT1-mediated internalization of this complex.

Although the results presented above may suggest that the sodium-independent transporter CAT1 is implicated in the transport of **Tc2** across cell membranes, the contribution of sodium-dependent transporters such as amino acid transport systems $b^{0,+}AT$ and $ATB^{0,+}$ cannot be discarded at this point.

The driving force for **Tc2** internalization could also be strengthened by the simultaneous contribution of a specific mechanism of retention in the cytosol of the cell, namely by interaction of the complex with cytosolic NOS, which we have already demonstrated to be the case for complexes of a related family mentioned above.²² To test this hypothesis we have studied the ability of **Re1** and **Re2**, cold surrogates of **Tc1** and **Tc2**, to be recognized by NOS both in enzymatic assays with purified iNOS and in LPS-activated macrophages. For the sake of comparison, the corresponding ligands **L1** and **L2** were also tested.

Firstly, the molecules were tested as NO-producing substrates using mouse recombinant iNOS. The iNOS activity was determined spectrophotometrically by monitoring the NO-mediated conversion of oxyhemoglobin to methemoglobin at 401 nm and 421 nm as previously described.^{22, 29} Table 1 displays the kinetic parameter K_m for all compounds, which was determined by the method of Eisenthal and Cornish-Bowden.

Table 1. K_m values for L-Arg, **L1**, **L2**, **Re1** and **Re2**

| Compound | K_m values/ μM |
|--------------|-----------------------------|
| L-Arg | 3.00 ± 1.00 |
| L1 | 50 |
| Re1 | 1093 |
| L2 | 1200 |
| Re2 | > 2500 |

The ability of **L1** ($K_m = 50 \mu\text{M}$) to interact specifically with the active site of iNOS, leading to NO production, is considerably higher than that observed for **L2** ($K_m = 1200 \mu\text{M}$) however, both compounds are poorer substrates than the endogenous substrate (L-Arg) of the enzyme ($K_m = 3 \mu\text{M}$). Metallation of **L1** and **L2** led to complexes with even lower NO-producing properties, as evidenced by the determined K_m values (**Re1**, $K_m = 1093 \mu\text{M}$; **Re2**, $K_m > 2500 \mu\text{M}$).

The effect of the same compounds was also studied in RAW 264.7 macrophages after treatment with lipopolysaccharide (LPS), which leads to increased NO biosynthesis due to iNOS overexpression.^{22, 34} This cellular model is very useful, as it allows to assess both the capacity of the compounds to cross cellular membranes, a key feature for improving tracer uptake *in vivo*, and the intracellular interaction with iNOS enzyme via the quantitation of the NO release.

The results, which represent the ability of the compounds to be recognized and used as NOS substrates and, consequently, their efficacy in NO biosynthesis in LPS-activated macrophages cultured in arginine-free medium, are presented in Figure 6.




Figure 6: Effect of **L1**, **L2**, **Re1** and **Re2** on NO biosynthesis by LPS-induced RAW 264.7 macrophages. Data are expressed as % of nitrite accumulation of the L-Arg control (mean \pm S.D. -standard deviation, n = 8). The experiment was repeated three times with comparable results

When LPS-induced macrophages were incubated in L-Arg-free culture medium, negligible NO production was observed. However, when macrophages were incubated with various concentrations of L-Arg (100 – 700 μ M) a high nitrite accumulation in the extracellular medium was observed (data not shown).

Incubation of LPS-treated macrophages with the new compounds showed that **L¹** and **L²** are recognized as substrates by the enzyme, confirmed by NO production (ca. 50%). When the corresponding rhenium compounds **Re1** and **Re2** are incubated with the cells the production of NO was negligible. These results are consistent with those obtained in the enzymatic assays, where it has been possible to conclude that metallation of conjugates **L1** and **L2** led to decreased affinities towards the enzyme.

We also performed a parallel viability assay to assess the intrinsic cytotoxicity of the compounds at the concentration used in the NO assay. The compounds were tested in the

absence or presence of L-Arg (Figure 7) in order to evaluate whether the viability is related with the presence/absence of NO, a key signaling mediator in several metabolic processes.




Figure 7- Viability in macrophages RAW 264.7 incubated with the compounds, in the absence or presence of L-arginine in the culture medium. Viability of cells incubated with LPS and L-arginine was considered as 100 % (mean \pm S.D., n = 8). This experiment was repeated two/three times with comparable results.

In the case of **L¹** and **L²** the viability is similar in both conditions (approximately 100%). For metal complexes **Re1** and **Re2**, which are not utilized as substrates, the viability is decreased in the absence of L-Arg (ca. 50%), most likely determined by impaired NO biosynthesis, with a value comparable to the control performed in the absence of that amino acid (ca. 50%). However, the viability of the metal complexes is comparable to the control when that amino acid is present in the culture medium, showing that the compounds by themselves are not toxic at the concentrations tested. All together, these results indicated that viability was being determined by NO biosynthesis.

Conclusions

We have prepared and characterized new conjugates comprising a pyrazolyl-diamine chelating unit and a pendant L-Arg moiety linked by a propyl (L^1) or a hexyl linker (L^2). The conjugates reacted with the organometallic precursors $fac-[M(CO)_3(H_2O)_3]^+$ ($M = {}^{99m}\text{Tc}/\text{Re}$) yielding the radioactive complexes of the type $fac-[{}^{99m}\text{Tc}(\text{CO})_3(\kappa^3\text{-L})]$ (**Tc1**, $L = L^1$; **Tc2**, $L = L^2$) and the respective “cold” surrogates **Re1** and **Re2**. **Tc2** exhibited moderate levels of time-dependent internalization in three different human tumoral cell models, with about 3% of the applied activity internalized after 4h at 37 °C, corresponding to 21% of the total cell-associated activity. Preliminary mechanistic studies suggest that internalization of **Tc2** is mediated by cationic amino acid transporters, namely system γ^+ , although more extensive amino acid transport assays will be needed to fully address that issue in the cancer lines studied. In addition, enzymatic assays with purified iNOS and studies with LPS-activated macrophages demonstrate that the surrogate complex **Re2** does not recognize a putative target of **Tc2** in the cytosol. Nevertheless, **Tc2** is a rare example of a metal complex whose entrance into cells seems to be mediated by cationic amino acid transporters. This finding might open the way towards the development of novel families of metal-based radiotracers to probe metabolically active cancer cells.

Experimental Section

General procedures and materials: All chemicals and solvents were of reagent grade and were used without purification unless stated otherwise. The BOC-protected precursor *tert*-butyl 2-((3-aminopropyl) (2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl) amino) ethylcarbamate (**Pz-C₃NH₂(Boc)**), *tert*-butyl 2-((6-aminoethyl)(2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl)amino)ethylcarbamate (**Pz-C₆NH₂(Boc)**), $[\text{Re}(\text{CO})_3(\text{H}_2\text{O})_3]\text{Br}$ were prepared according to published methods.^{22, 35} 2-(*tert*-Butoxycarbonylamino)-5-guanidinopentanoic acid (*N*- α -Boc-L-Arg) was purchased from Sigma Aldrich as well as all other chemicals not specified above. $\text{Na}[{}^{99m}\text{TcO}_4]$ was eluted from a

$^{99}\text{Mo}/^{99\text{m}}\text{Tc}$ generator, using 0.9 % saline. The radioactive precursor $\text{fac-}[^{99\text{m}}\text{Tc}(\text{CO})_3(\text{H}_2\text{O})_3]^+$ was prepared using a IsoLink[®] kit (Covidean Malinckrodt, Inc.). ^1H and ^{13}C NMR spectra were recorded at room temperature on a Varian Unity 300 MHz spectrometer. ^1H and ^{13}C chemical shifts were referenced with the residual solvent resonances relatively to tetramethylsilane. The spectra were assigned with the help of 2D experiments (^1H - ^1H correlation spectroscopy, COSY and ^1H - ^{13}C heteronuclear single quantum coherence, HSQC). Assignments of the ^1H and ^{13}C NMR resonances are given in accordance with the identification system shown in Scheme 1. Infrared spectra were recorded as KBr pellets on a Bruker Tensor 27 spectrometer. All compounds were characterized by electrospray ionization mass spectrometry (ESI-MS) using a Bruker model Esquire 3000 plus. HPLC analyses were performed on a Perkin Elmer LC pump 200 coupled to a Shimadzu SPD 10AV UV/Vis and to a Berthold-LB 509 radiometric detector, using an analytic Macherey-Nagel C18 reversed-phase column (Nucleosil 100-5, 250 x 3 mm) with a flow rate of 0.5 mL/min. Eluents: aqueous 0.1 % $\text{CF}_3\text{CO}_2\text{H}/\text{MeOH}$. Gradient: $t = 0\text{-}5$ min: 10 % MeOH; $5\text{-}30$ min: $10 \rightarrow 100$ % MeOH; $30\text{-}34$ min: 100 % MeOH; $34\text{-}35$ min: $100 \rightarrow 10$ % MeOH; $35\text{-}40$ min: 10 % MeOH.

Purification of the inactive compounds was achieved on a semi-preparative Macherey-Nagel C18 reversed-phase column (Nucleosil 100-7, 250 x 8 mm) or on a preparative Waters $\mu\text{Bondapak}$ C18 (150 x 19 mm) with a flow rate of 2.0 mL/min and 5.5 mL/min, respectively. UV detection: 220 or 254 nm. Eluents: aqueous 0.1 % $\text{CF}_3\text{CO}_2\text{H}/\text{MeOH}$. Gradient: $t = 0\text{-}5$ min: 10 % MeOH; $5\text{-}30$ min: $10 \rightarrow 100$ % MeOH; $30\text{-}34$ min: 100 % MeOH; $34\text{-}35$ min: $100 \rightarrow 10$ % MeOH; $35\text{-}40$ min: 10 % MeOH.

Aliquots of ~ 5 mg of pure compounds (≥ 98 % ascertained by RP-HPLC) were lyophilized in microcentrifuge tubes and used for radioactive labelling and *in vitro* studies.

Synthesis of *tert*-butyl 2-((3-(2-amino-5-guanidinopentanamido)propyl)(2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl)amino)ethylcarbamate (L¹-Boc)

To a solution of **Pz-C₃NH₂(Boc)** (0.050 g, 0.140 mmol) in dimethylformamide (DMF) were added triethylamine (0.036 g, 0.365 mmol) and O-benzotriazol-1-yl-N,N,N',N'-tetramethyluronium hexafluorophosphate (HBTU, 0.054 g, 0.145 mmol). After 10 min, N- α -Boc-L-Arg (0.039 g, 0.142 mmol) was added, and the reaction mixture stirred at room temperature under a nitrogen atmosphere for 2 h. The solvent was removed under vacuum, and the residue purified by silica gel column chromatography using a gradient of MeOH (0 \rightarrow 100 %) in CHCl₃. The intermediate **L¹-Boc** was obtained as a yellowish oil. Yield: 58.1 % (0.050 g, 0.082 mmol).

¹H-NMR (300 MHz, CDCl₃): δ_H (ppm) 7.69 (2H, br s, NH), 6.48 (1H, br s, NH), 5.70 (1H, s, CH^b), 5.10 (1H, br s, NH), 5.01 (2H, t, CH₂^d), 4.57 (1H, br s, NH), 3.35 (2H, t, CH₂^e), 3.07 (1H, br m, CH^k), 2.86-2.80 (4H, m, CH₂^{g,j}), 2.35 (2H, t, CH₂ⁿ), 2.32 (2H, t, CH₂^f), 2.23 (2H, t, CH₂^g), 2.19 (3H, s, CH₃^{Pz}), 2.16 (3H, s, CH₃^{Pz}), 1.40-1.38 (18H, s, CH₃^{Boc}), 1.39 – 1.15 (6H, m, CH₂^{i,l,m}).

¹³C-NMR (75.5 MHz, CDCl₃): δ_C (ppm) 174.3 (CO), 172.6 (CO), 159.4 (C^o), 147.3 (C^{Pz}), 143.9 (C^{Pz}), 106.9 (C^{Pz}), 79.9 (C(CH₃)₃), 56.7 (C^e), 54.1 (C^k), 53.8 (C^f), 51.6 (C^h), 49.2 (C^d), 41.2 (Cⁿ), 38.9 (C^g), 38.0 (C^j), 30.8 (Cⁱ), 28.8 (C(CH₃)₃), 25.4 (C^l), 24.6 (C^m), 13.4 (CH₃Pz), 11.0 (CH₃Pz).

Synthesis of *tert*-butyl 2-((6-(2-amino-5-guanidinopentanamido)hexyl)(2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl)amino)ethylcarbamate (L²-Boc)

To a solution of **Pz-C₆NH₂(Boc)** (0.050 g, 0.131 mmol) in DMF were added triethylamine (0.036 g, 0.365 mmol) and O-benzotriazol-1-yl-N,N,N',N'-tetramethyluronium hexafluorophosphate (HBTU, 0.054 g, 0.145 mmol). After 10 minutes, N- α -Boc-L-Arg (0.039 g, 0.142 mmol) was added, and the reaction mixture stirred at room temperature under a nitrogen atmosphere for 2 h. The solvent was then removed under vacuum, and the residue purified by silica gel column chromatography using a gradient of MeOH (0 \rightarrow 100 %) in CHCl₃. The intermediate **L²-Boc** was obtained as a yellowish oil. Yield: 60.2 % (0.050 g, 0.078 mmol).

¹H-NMR (300 MHz, CDCl₃): δ_H (ppm) 7.70 (2H, br s, NH), 6.51 (1H, br s, NH), 5.73 (1H, s, CH^b), 5.14 (1H, br s, NH), 5.05 (2H, t, CH₂^d), 4.61 (1H, br s, NH), 3.37 (2H, t, CH₂^e), 3.13 (1H, br m, CHⁿ), 3.13-3.06 (4H, m, CH₂^{g,m}), 2.65 (2H, t, CH₂^h), 2.52 (2H, t, CH₂^q), 2.43 (2H, t, CH₂^f), 2.20 (3H, s, CH₃^{Pz}), 2.18 (3H, s, CH₃^{Pz}), 1.80 (2H, q, C^o), 1.80 (2H, m, C^l), 1.42-1.40 (18H, s, CH₃^{Boc}), 1.32 – 1.26 (6H, m, CH₂^{ij,k}).

¹³C-NMR (75.5 MHz, CDCl₃): δ_c (ppm) 174.5 (CO), 172.8 (CO), 158.4 (C^r), 147.5 (C^{Pz}), 144.2 (C^{Pz}), 105.9 (C^{Pz}), 79.5 (C(CH₃)₃), 56.7 (C^e), 56.3 (C^h), 54.1 (C^f), 53.6 (Cⁿ), 49.4 (C^d), 41.5 (C^q), 39.3 (C^m), 38.8 (C^g), 30.6 (Cⁿ), 28.9 (C(CH₃)₃), 28.3 (Cⁱ), 27.6 (C^j), 26.4 (C^k), 25.6 (C^o), 24.9 (C^p), 13.5 (CH₃^{Pz}), 11.3 (CH₃^{Pz}).

Synthesis of 2-amino-N-(3-((2-aminoethyl)(2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl)amino)-propyl)-5-guanidinopentanamide (L¹)

Compound **L¹** was obtained directly by dissolving **L¹-Boc** (0.050 g, 0.082 mmol) in a mixture CH₂Cl₂–TFA (1 mL – 3 mL) and allowed to react for 3 h at room temperature with stirring. The residue obtained after evaporation of the solvents was dissolved in water, filtered through a 0.45 μm Millipore filter, and purified by preparative RP-HPLC. The fractions containing **L¹** were collected and the solvent removed to provide a clear viscous oil. Yield: 78 % (0.026 g, 0.061 mmol, calcd. for C₁₉H₄₁N₉O).

¹H-NMR (300 MHz, D₂O): δ_H (ppm) 5.99 (1H, s, H^b), 4.37 (2H, s, H^d), 3.86 (1H, t, Hⁿ), 3.49 (2H, m, H^e), 3.39 (2H, m, H^f), 3.32 (2H, t, H^g), 3.24-3.06 (6H, t, H^{q, h, j}), 2.19 (3H, s, CH₃^{Pz}), 2.10 (3H, s, CH₃^{Pz}), 1.81 (4H, m, H^{o, i}), 1.53 (2H, m, H^p).

¹³C-NMR (75.5 MHz, D₂O): δ_c (ppm) 169.9 (CO), 156.9 (C^r), 148.9 (C^c), 144.4 (C^a), 107.3 (C^b), 52.9 (Cⁿ), 51.8 (C^h), 51.7 (C^e), 49.9 (C^f), 42.2 (C^d), 40.4 (C^q), 36.4 (C^j), 33.9 (C^g), 28.1 (Cⁱ), 23.8 (C^o), 23.4 (C^p), 11.5 (CH₃^{Pz}), 10.1 (CH₃^{Pz}).

RP-HPLC (t_R): 18.7 min.

ESI-MS (+) (m/z): 412.7 [M + H]⁺, calcd. for C₁₉H₄₁N₉O = 411.6.

IR (KBr, cm⁻¹): 3445 M v(NH₂, NH); 1612 S v(C=O), δ(NH₂), δ(NH, amide), 1480 w v(CN, amide); 1220 e 1137 S v(C-N); 909 w, 836 w, 765 w.

Synthesis of 2-amino-N-(6-((2-aminoethyl)(2-(3,5-dimethyl-1H-pyrazol-1-yl)ethyl)amino)-hexyl)-5-guanidinopentanamide (L²)

Removal of Boc protecting group of L²-Boc was done following the methodology described for L¹-Boc. The residue obtained after evaporation of the solvents was dissolved in water, filtered through a 0.45 μm Millipore filter, and purified by preparative RP-HPLC. The fractions containing L² were collected and the solvent removed to provide a yellow viscous oil. Yield: 90.4 % (0.019 g, 0.043 mmol, calcd. for C₂₁H₄₃N₉O).

¹H-NMR (300 MHz, D₂O): δ_H (ppm) 6.02 (1H, s, H^b), 4.42 (2H, t, H^d), 3.83 (1H, t, Hⁿ), 3.69 (2H, t, H^e), 3.51 (2H, m, H^f), 3.35 (2H, m, H^g, H^{g'}), 3.13 (6H, m, H^{q, h, m}), 2.21 (3H, t, CH₃^{Pz}), 2.13 (3H, s, CH₃^{Pz}), 1.78 (2H, m, H^o), 1.55-1.49 (4H, m, H^{i, p}), 1.42-1.32 (2H, m, H^l), 1.22 (4H, m, H^{j, k}).

¹³C-NMR (75.5 MHz, D₂O): δ_c (ppm) 169.6 (CO), 156.8 (C^r), 148.2 (C^{Pz}), 145.8 (C^{Pz}), 107.9 (C^b), 53.9 (C^q), 52.9 (Cⁿ), 51.0 (C^e), 49.7 (C^f), 42.0 (C^d), 40.3 (C^m), 39.4 (C^h), 33.6 (C^g), 28.0 (C^{o, l}), 25.6 (C^p), 25.2 (Cⁱ), 23.7 (C^k), 22.8 (C^j), 10.9 (CH₃^{Pz}), 10.1 (CH₃^{Pz}).

RP-HPLC (t_R): 19.5 min.

ESI-MS (+) (m/z): 220 [M + H]²⁺, 439 [M+H]⁺, calcd. for C₂₁H₄₃N₉O = 438.

IR (KBr, cm⁻¹): 3312 M v(NH₂, NH); 1671 S v(C=O), δ(NH₂), δ(NH, amide), 1421 w v(CN, amide); 1209 and 1190 S v(C-N); 898 w, 823 w, 748 w.

General procedure for the preparation of the Re complexes *fac*-[Re(CO)₃(k³-L)] (Re1, L = L1; Re2, L = L2)

[Re(CO)₃(H₂O)₃]Br was reacted with equimolar amounts of **L**¹ or **L**² in refluxing water for 12 h. The solvent was concentrated to ¼ of the volume and the resulting solution was centrifuged and the supernatant purified by preparative RP-HPLC.

Synthesis of fac-[Re(CO)₃(k³-L¹)]⁺ (Re1): Starting from 0.020 g (0.048 mmol) of **L**¹, a colorless oil formulated as **Re1** was obtained. Yield: 33.3 % (0.010 g, 0.014 mmol, calcd. for C₂₂H₄₁N₉O₄Re).

¹H-NMR (300 MHz, D₂O): δ_H (ppm) 6.08 (1H, s, H^b), 5.10 (1H, s, NH), 4.35 (1H, dd, H^d), 4.14-3.98 (1H, m, H^{d''}), 3.87 (1H, t, Hⁿ), 3.71-3.66 (2H, m, NH), 3.59-3.50 (1H, m, CH^h), 3.40-3.19 (4H, m, H^{h', e', j}₂), 3.11 (3H, m, H^{g', q}₂), 2.75 (2H, m, H^f), 2.67-2.63 (1H, m, H^{e''}), 2.50-2.39 (1H, m, H^{g''}), 2.30 (3H, s, CH₃^{Pz}), 2.19 (3H, s, CH₃^{Pz}), 2.10-1.91 (1H, m, Hⁱ), 1.86-1.75 (3H, m, H^{i', o}₂), 1.61-1.50 (2H, m, H^p₂).

¹³C-NMR (75.5 MHz, D₂O): δ_C (ppm) 196.1, 194.9, 194.7 (CO, Re), 171.6 (C=O), 164.7 (Cⁱ), 155.7 (C^{Pz}), 146.2 (C^{Pz}), 109.8 (C^{Pz}), 66.7 (C^h), 63.7 (C^f), 55.1 (Cⁿ) 55.4 (C^e), 49.0 (C^d), 44.1 (C^g), 42.4 (C^j), 39.3 (C^q), 30.1 (C^o), 26.2 (Cⁱ), 25.8 (C^p), 17.2 (CH₃^{Pz}), 12.8 (CH₃^{Pz}).

RP-HPLC (t_R): 25.1 min

ESI-MS (+) (m/z): 342 [M+2H]²⁺, calcd. for C₂₁H₃₇N₉O₄Re = 681.3.

IR (KBr, cm⁻¹): 3276 M ν(NH₂ NH); 2026, 1915 S (CO); 1676 S (C=O), δ(NH₂), 1208 – 1135 M ν(CN); 876 w; 758 w; 683 w.

Synthesis of fac-[Re(CO)₃(k³-L²)]⁺ (Re2): Starting from 0.020 g (0.045 mmol) of **L**², a colorless oil formulated as **Re**² was obtained. Yield: 37.5 % (0.012 g, 0.017 mmol, calcd. for C₂₄H₄₃N₉O₄Re).

¹H-NMR (300 MHz, D₂O): δ_H (ppm) 6.05 (1H, s, H^b), 5.04 (1H, s, NH), 4.32 (1H, dd, H^d), 4.08 (1H, dd, H^{d''}), 3.82 (1H, t, Hⁿ), 3.61 (2H, s, NH), 3.50-3.40 (1H, m, H^h), 3.30-3.18 (3H, m, H^{h', m}₂), 3.07

(3H, m, H^{e', g', q}), 2.70 (2H, m, H^f), 2.53 (1H, m, H^{e''}), 2.40 (1H, m, H^{g''}), 2.27 (3H, s, H^{Pz}), 2.16 (3H, s, H^{Pz}), 1.76 (3H, m, H^{o', o'', i''}), 1.69-1.58 (1H, m, H^{i''}), 1.49 (4H, m, H^{p, l}) 1.61-1.50 (2H, m, H^{j, k}).

¹³C-NMR (75.5 MHz, D₂O): δ_c (ppm) 194.7, 194.4, 193.0 (CO, Re); 169.2 (C=O), 156.9 (C^f), 153.7 (C^{Pz}), 144.2 (C^{Pz}), 107.9 (C^{Pz}), 67.6 (C^h), 61.9 (C^f), 53.3 (C^e), 53.3 (Cⁿ), 47.0 (C^d), 42.2 (C^g), 40.5 (C^m), 39.6 (C^q), 28.2 (C^l), 28.2 (C^o), 25.9 (C^{j, k}), 24.4 (Cⁱ), 23.8 (C^l), 15.3 (CH₃^{Pz}), 10.9 (CH₃^{Pz}).

RP-HPLC (t_R): 25.5 min.

ESI-MS (+) (m/z): 355 [M + 2H]²⁺, calcd. for C₂₄H₄₃N₉O₄Re = 708.

IR (KBr, cm⁻¹): 3274 M ν (NH₂ NH); 2028, 1914 S (CO); 1675 S (C=O), δ (NH₂), 1206 – 1135 M ν (CN); 877 w; 760 w; 684 w.

General method for the synthesis of the ^{99m}Tc(I) complexes *fac*-[^{99m}Tc(CO)₃(k³-L)] (L = L¹ and L²)

In a nitrogen-purged glass vial, 100 mL of a 10⁻⁴ M aqueous solution of the compounds **L1** or **L2** were added to 900 mL of a solution of the organometallic precursor *fac*-[^{99m}Tc(CO)₃(H₂O)₃]⁺ (1 – 2 mCi) in saline or phosphate buffer pH 7.4. The reaction mixture was then heated to 100 °C for 30min, cooled on an ice bath and the final solution analyzed by RP-HPLC. Retention times: 25.9 min (**Tc1**), 26.6 min (**Tc2**). Complexes **Tc3** - **Tc5** were prepared and characterized as described previously.^{22, 28}

In vitro stability studies in plasma

100 μ L of **Tc1** or **Tc2** were added to 500 μ L of human plasma and incubated at 37 °C. After 24 h, aliquots (100 μ L) were taken and the plasmatic proteins precipitated with ethanol (200 μ L). The plasma was centrifuged at 3000 rpm for 15 min at 4 °C and the supernatant (protein-free plasma) filtered through a Millipore filter (0.22 μ m), and analyzed by RP-HPLC.

Enzymatic assays

The iNOS activity assay was based on the method of hemoglobin assay previously described by Hevel and Marletta with slight modifications.^{36, 37} The kinetics parameters for iNOS were determined using initial rate analysis. Initial rate data were fitted to irreversible single substrate Michaelis–Menten models. The kinetic parameters were determined using the direct linear plot of Eisenthal and Cornish-Bowden and the Hyper software (J.S. Easterby, University of Liverpool, UK; <http://www.liv.ac.uk/~jse/software.html>).³⁸ This method was chosen primarily because of its robustness.³⁹ The K_m values represent a mean of triplicate measurements. Standard deviations of ± 5 to 10 % were observed.

Preparation of oxyhemoglobin

Oxyhemoglobin was prepared using a previously described protocol with some modifications.⁴⁰ Briefly, bovine hemoglobin in 50 mM HEPES pH 7.4 was reduced to oxyhemoglobin with 10-fold molar excess of sodium dithionite. The sodium dithionite was later removed by dialysis against 50 volumes of HEPES buffer for 18 h at 4 °C. The buffer was replaced 3 times. The concentration of oxyhemoglobin was determined spectrophotometrically using $\epsilon_{415\text{ nm}} = 131\text{ mM}^{-1}\text{ cm}^{-1}$. Oxyhemoglobin was stored at - 80 °C before use.

Determination of K_m values

All initial velocity measurements were recorded at 37 °C. Total reaction volumes were 1500 μL and contained 50 mM HEPES pH 7.4, 6 mM oxyhemoglobin, 200 mM NADPH, 10 mM Tetrahydrobiopterin (BH_4), 100 mM DTT and increasing concentrations of L-Arg, **L1**, **L2**, **Re1** and **Re2** (20 – 500 mM). Magnetic stirring in the spectrophotometer cuvette was essential to maintain isotropic conditions. Reactions were initiated by the addition of iNOS enzyme ($\sim 1\text{ U}$) to the pre warmed cuvette ($\sim 5\text{ min}$). The NO-mediated conversion of oxyhemoglobin to methemoglobin was followed by monitoring the increase in absorbance at dual wavelength (401

and 421 nm) for 10 min.⁴¹ Controls were performed in the same conditions without iNOS enzyme.

Cell culture

RAW 264.7 macrophages, and the following human tumoral cell lines: HeLa cervical cancer, A375 melanoma, MDA-MB-231 breast cancer and PC3 prostate cancer, were grown in Dulbecco's Modified Eagle Medium (DMEM) with GlutaMax I supplemented with 10% heat-inactivated fetal bovine serum (FBS) and 1% penicillin/streptomycin antibiotic solution (all from Invitrogen, UK). Cells were cultured in a humidified atmosphere of 95 % air and 5 % CO₂ at 37 °C, with the medium changed every other day.

Cellular uptake and internalization

Cellular uptake assays with **Tc1** - **Tc5** were performed in HeLa, A375, MDA-MB-231 and PC3 cell lines seeded at a density of 0.2 million cells/well in a 24-well tissue culture plates. Cells were allowed to attach overnight. On the following day cells were exposed to complexes (about 200,000 cpm in 0.5 mL of assay medium: Modified Eagle's Medium with 25 mM HEPES and 0.2% BSA) for a period of 5 min to 4 h. Incubation was terminated by removing the Tc-complexes and by washing cells twice with ice-cold PBS with 0.2% BSA. Then, cells were lysed by 10 min incubation with 1 M NaOH at 37°C and the activity of lysates were measured in a γ -counter. The percentage of cell-associated radioactivity was calculated and represented as a function of incubation time. Uptake studies were carried out using at least four replicates for each time point.

The cellular uptake of **Tc2** was also evaluated in the presence of N-ethylmaleimide (NEM), a specific inhibitor of system γ^+ . Cells were incubated with solutions of **Tc2** containing NEM (5 mM) during different periods (5, 15, 30 and 60 min) at 37 °C. The general procedure of cellular

uptake was followed. The inhibition of cellular uptake was expressed in percentage of the uptake of **Tc2** in the absence of inhibitor.

Internalization assays of **Tc2** complex were performed in HeLa, A375 and MDA-MB-231 cell lines seeded at a density of 0.2 million cells/well in a 24-well tissue culture plates. On the following day, cells were exposed to **Tc2** (about 200,000 cpm in 0.5 mL of assay medium) for a period of 5 min to 4h. Incubation was terminated by washing the cells with ice-cold assay medium. Cell surface-bound **Tc2** was removed by two steps of acid wash (50 mM glycine-HCl/100 mM NaCl, pH 2.8) at room temperature for 4 min. The pH was neutralized with cold PBS with 0.2% BSA. The cells were then lysed by 10 min incubation with 0.5 N NaOH at 37 °C to determine internalized **Tc2**. The activity of the lysates and cell surface-bound fractions were measured in a γ -counter.

Internalization assays for **Tc2** were also performed in the presence of L-Lysine. For this study, A375, HeLa and MDA-231 cells were incubated 2h at 37°C with solutions of **Tc2** containing different concentrations of L-Lysine (0, 0.5, 1, 5 and 10mM).

Cell viability determination

Cell viability was evaluated by using a colorimetric method based on the tetrazolium salt [3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT), which is reduced by viable cells to yield purple formazan crystals.⁴² RAW 264.7 macrophages in DMEM medium without arginine supplemented with 10% FBS were seeded in 96-well plates at a density of 9×10^4 cells per well, immediately induced with LPS (2 mg/mL) for 4 h, and then incubated for 24 h in the presence of the compounds (500 μ M) and arginine (when indicated).²² At the end of the incubation period the media was removed and the cells were incubated with MTT (0.5 mg/mL in culture medium; 200 μ L) for 3 - 4 h at 37 °C and 5 % CO₂. The purple formazan crystals formed inside the cells were then dissolved in 200 μ L of DMSO by thorough shaking, and the absorbance was read at

570 nm, using a plate spectrophotometer (Power Wave Xs; Bio-Tek). Each test was performed with at least six replicates and repeated at least 2 times. The result was expressed as percentage of the surviving cells in relation with the control.

Assay of iNOS activity in vivo

RAW 264.7 macrophages in DMEM medium without arginine supplemented with 10% FBS DMEM, were plated at a density of 9×10^4 cells per well in 96-well plates. Cells were immediately induced with 10 μ L of LPS (2 μ g/mL in PBS) for 4 h, and then incubated for 24 h in the presence of the compounds (500 μ M). At the end of the incubation period, the culture medium was collected and assayed for nitrite production using the commercially available Griess reagent (1 % sulfanilamide, 0.1 % *N*-1-naphthyl ethylenediamine, 2.5 % orthophosphoric acid; Sigma-Aldrich). Briefly, 50 μ L of Griess reagent was mixed with an equal volume of medium at room temperature and absorbance was measured at 540 nm after 10 min. Fresh culture medium served as the blank in all experiments. Each experiment was performed with six replicates and repeated three times.

Western blot

Western blot experiments were performed to demonstrate the expression of CAT-1 in the four human tumoral cell lines (A375, HeLa, MDA-MB-231 and PC3). Cells were lysed with CellLyctic™ M Extraction Reagent (Sigma) supplemented with Complete Protease Inhibitor Cocktail tablets (Roche). After 15 min at 4° C with regular shaking, lysates were collected and centrifuged at 14000 g for 10 min to pellet the cellular debris, and the supernatants removed for further use. The total protein content was determined using the DC Protein Assay Kit (BioRad). Samples (50 μ g of protein) were separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and transferred onto nitrocellulose membranes. Blots were blocked with 5% nonfat dry milk in PBS-T for 2 h and then incubated for 1 h with the primary antibodies against

CAT-1 (1:200, Santa Cruz, sc-66824) and actin (1:15000, Sigma, A3853). Membranes were washed with PBS-T and incubated for 1 h with the secondary antibodies in a 1:3000 dilution (goat anti-rabbit IgG-HRP for α -CAT-1 and goat anti-mouse IgG-HRP for α -actin, BioRad). Membranes were developed using the SuperSignal West Pico Substrate kit (Pierce, Rockford, IL) according to the manufacturer's instructions.

Acknowledgments

This work has been partially supported by the *Fundação para a Ciência e Tecnologia* (FCT), Portugal, through the UID/Multi/04349/2013 project. V. F. C. F. thanks FCT for a PhD grant (SFRH/BD/108623/2015). Dr. J. Marçalo is acknowledged for performing the ESI-MS analyses. The QITMS instrument was acquired with the support of Contract REDE/1503/REM/2005 - ITN of FCT and is part of RNEM.

References

1. S. L. Pimlott and A. Sutherland, *Chemical Society Reviews*, 2011, **40**, 149-162.
2. S. N. Histed, M. L. Lindenberg, E. Mena, B. Turkbey, P. L. Choyke and K. A. Kurdziel, *Nuclear Medicine Communications*, 2012, **33**, 349-361.
3. H. Schoder and S. M. Larson, *Seminars in Nuclear Medicine*, 2004, **34**, 274-292.
4. A. Sundin, U. Garske and H. Orlefors, *Best Practice & Research Clinical Endocrinology & Metabolism*, 2007, **21**, 69-85.
5. J. McConathy and M. M. Goodman, *Cancer and Metastasis Reviews*, 2008, **27**, 555-573.
6. J. McConathy, W. Yu, N. Jarkas, W. Seo, D. M. Schuster and M. M. Goodman, *Medicinal Research Reviews*, 2012, **32**, 868-905.
7. R. Hyde, P. M. Taylor and H. S. Hundal, *Biochemical Journal*, 2003, **373**, 1-18.
8. E. I. Closs, J. P. Boissel, A. Habermeier and A. Rotmann, *Journal of Membrane Biology*, 2006, **213**, 67-77.
9. C. Huang and J. McConathy, *Journal of Nuclear Medicine*, 2013, **54**, 1007-1010.
10. K. K. S. Sai, C. Huang, L. Yuan, D. Zhou, D. Piwnica-Worms, J. R. Garbow, J. A. Engelbach, R. Mach, K. M. Rich and J. McConathy, *Journal of Nuclear Medicine*, 2013, **54**, 1120-1126.
11. E. I. Closs, A. Simon, N. Vekony and A. Rotmann, *Journal of Nutrition*, 2004, **134**, 2752S-2759S.
12. S. Moncada, R. M. J. Palmer and E. A. Higgs, *Pharmacological Reviews*, 1991, **43**, 109-142.
13. D. Fukumura, S. Kashiwagi and R. K. Jain, *Nature Reviews Cancer*, 2006, **6**, 521-534.
14. P. Mukherjee, M. A. Cinelli, S. Kang and R. B. Silverman, *Chemical Society Reviews*, 2014, **43**, 6814-6838.

15. M. M. Phillips, M. T. Sheaff and P. W. Szlosarek, *Cancer Research and Treatment*, 2013, **45**, 251-262.
16. N. Gupta, S. Miyauchi, R. G. Martindale, A. V. Herdman, R. Podolsky, K. Miyake, S. Mager, P. D. Prasad, M. E. Ganapathy and V. Ganapathy, *Biochimica Et Biophysica Acta-Molecular Basis of Disease*, 2005, **1741**, 215-223.
17. N. Gupta, P. D. Prasad, S. Ghamande, P. Moore-Martin, A. V. Herdman, R. G. Martindale, R. Podolsky, S. Mager, M. E. Ganapathy and V. Ganapathy, *Gynecologic Oncology*, 2006, **100**, 8-13.
18. S. Karunakaran, S. Ramachandran, V. Coothankandaswamy, S. Elangovan, E. Babu, S. Periyasamy-Thandavan, A. Gurav, J. P. Gnanaprakasam, N. Singh, P. V. Schoenlein, P. D. Prasad, M. Thangaraju and V. Ganapathy, *Journal of Biological Chemistry*, 2011, **286**, 31830-31838.
19. Y. Lu, W. Wang, J. Wang, C. Yang, H. Mao, X. Fu, Y. Wu, J. Cai, J. Han, Z. Xu, Z. Zhuang, Z. Liu, H. Hu and B. Chen, *Plos One*, 2013, **8**.
20. S. A. Abdelmagid, J. A. Rickard, W. J. McDonald, L. N. Thomas and C. K. L. Too, *Journal of Cellular Biochemistry*, 2011, **112**, 1084-1092.
21. Y. Liu, J. K. Pak, P. Schmutz, M. Bauwens, J. Mertens, H. Knight and R. Alberto, *Journal of the American Chemical Society*, 2006, **128**, 15996-15997.
22. B. L. Oliveira, P. D. Raposinho, F. Mendes, F. Figueira, I. Santos, A. Ferreira, C. Cordeiro, A. P. Freire and J. D. G. Correia, *Bioconjugate Chemistry*, 2010, **21**, 2168-2172.
23. A. de Almeida, B. L. Oliveira, J. D. G. Correia, G. Soveral and A. Casini, *Coordination Chemistry Reviews*, 2013, **257**, 2689-2704.
24. B. L. Oliveira, I. S. Moreira, P. A. Fernandes, M. J. Ramos, I. Santos and J. D. G. Correia, *Journal of Molecular Modeling*, 2013, **19**, 1537-1551.
25. B. L. Oliveira, I. S. Moreira, P. A. Fernandes, M. J. Ramos, I. Santos and J. o. D. G. Correia, *Journal of Molecular Graphics and Modelling*, 2013, **45**, 13-25.
26. B. L. Oliveira, M. Morais, F. Mendes, I. S. Moreira, C. Cordeiro, P. A. Fernandes, M. J. Ramos, R. Alberto, I. Santos and J. D. G. Correia, *Chemical Biology & Drug Design*, 2015, **86**, 1072-1086.
27. M. Morais, A. Paulo, L. Gano, I. Santos and J. D. G. Correia, *Journal of Organometallic Chemistry*, 2013, **744**, 125-139.
28. B. L. Oliveira, J. D. G. Correia, P. D. Raposinho, I. Santos, A. Ferreira, C. Cordeiro and A. P. Freire, *Dalton Transactions*, 2009, 152-162.
29. B. L. Oliveira, J. D. G. Correia, P. D. Raposinho, I. Santos, A. Ferreira, C. Cordeiro and A. P. Freire, *Dalton Transactions*, 2009, **0**, 152-162.
30. M. Pan, H. A. Choudry, M. J. Epler, Q. H. Meng, A. Karinch, C. M. Lin and W. Souba, *Journal of Nutrition*, 2004, **134**, 2826S-2829S.
31. D. Fotiadis, Y. Kanai and M. Palacin, *Molecular Aspects of Medicine*, 2013, **34**, 139-158.
32. C. Fleck, M. Schwertfeger and P. M. Taylor, *Amino Acids*, 2003, **24**, 347-374.
33. S. R. Beyer, R. T. Mallmann, I. Jaenecke, A. Habermeier, J. P. Boissel and E. I. Closs, *Journal of Biological Chemistry*, 2013, **288**, 30411-30419.
34. J. MacMicking, Q. W. Xie and C. Nathan, *Annual Review of Immunology*, 1997, **15**, 323-350.
35. N. Lazarova, S. James, J. Babich and J. Zubieta, *Inorganic Chemistry Communications*, 2004, **7**, 1023-1026.
36. J. Hevel and M. Marletta, *Methods Enzymol.*, 1994, **233**, 250 - 258.
37. S. Archer, *The FASEB Journal*, 1993, **7**, 349-360.
38. Robert Eisenthal and A. Cornish-Bowden, *Biochem J.*, 1974, **139**, 715 – 720.
39. A. Cornish-Bowden, *Fundamentals of Enzyme Kinetics*, Portland Press, London, 2nd edn., 1995.
40. D. J. Wolff, D. S. Gauld, M. J. Neulander and G. Southan, *Journal of Pharmacology and Experimental Therapeutics*, 1997, **283**, 265-273.

41. R. G. Knowles and S. Moncada, *Biochemical Journal*, 1994, **298**, 249-258.
42. T. Mosmann, *Journal of Immunological Methods*, 1983, **65**, 55-63.

Supplementary Information for

Technetium-99m complexes of L-arginine derivatives for targeting amino acid transporters

Maurício Morais,^{1,2,‡} Vera F. C. Ferreira,^{1,‡} Flávio Figueira,¹ Filipa Mendes,¹ Paula Raposinho,¹
Isabel Santos,¹ Bruno L. Oliveira,^{1,3} João D. G. Correia^{1,*}

¹Centro de Ciências e Tecnologias Nucleares, Instituto Superior Técnico, Universidade de Lisboa, Estrada Nacional 10 (km 139,7), 2695-066 Bobadela LRS, Portugal

²Current address: Division of Imaging Sciences and Biomedical Engineering, King's College London, 4th Floor Lambeth Wing, St Thomas' Hospital, London SE1 7EH, United Kingdom

³Current address: Department of Chemistry, University of Cambridge, Lensfield Road, Cambridge, United Kingdom

Submitted as a full article to: Dalton Trans., *Frontiers in Radionuclide Imaging and Therapy themed issue*

Keywords: Amino acid transporters, Arginine, Cancer, Imaging, Rhenium, Technetium

* Corresponding author: João D. G. Correia, Tel.: +351 21 994 62 33

E-mail address: jgalamba@ctn.tecnico.ulisboa.pt

‡ These authors contributed equally to the article.

RP-HPLC CHROMATOGRAMS





Figure S1. RP-HPLC chromatograms of **L¹** and **L²**. Macherey-Nagel C18 reversed-phase column (Nucleosil 100-5, 250 x 3 mm) with a flow rate of 0.5 mL/min. Eluents: aqueous 0.1 % CF₃CO₂H/MeOH. Gradient: t = 0-5 min: 10 % MeOH; 5-30 min: 10→100 % MeOH; 30-34 min: 100 % MeOH; 34-35 min: 100→10 % MeOH; 35-40 min: 10 % MeOH. Detection: $\lambda = 220$ nm.

NMR SPECTRA

¹H-NMR spectrum



¹³C-NMR spectrum





Figure S2. Assigned ¹H and ¹³C-NMR spectra of L¹.

¹H-NMR spectrum



¹³C-NMR spectrum




Figure S3. Assigned ¹H and ¹³C-NMR spectra of L².




Figure S4. RP-HPLC chromatograms of **Tc1** and **Tc2**. Macherey-Nagel C18 reversed-phase column (Nucleosil 100-5, 250 x 3 mm) with a flow rate of 0.5 mL/min. Eluents: aqueous 0.1 % $\text{CF}_3\text{CO}_2\text{H}$ /MeOH. Gradient: t = 0-5 min: 10 % MeOH; 5-30 min: 10 \rightarrow 100 % MeOH; 30-34 min: 100 % MeOH; 34-35 min: 100 \rightarrow 10 % MeOH; 35-40 min: 10 % MeOH. γ detection.




Figure S5. Stability of **Tc1** and **Tc2**. RP-HPLC chromatograms of **Tc1** and **Tc2** in phosphate buffered saline (PBS) pH 7.4 **(A)** and in PBS pH 7.4 with a 100 fold excess histidine **(B)** after 24 h at 37°C. Macherey-Nagel C18 reversed-phase column (Nucleosil 100-5, 250 x 3 mm) with a flow rate of 0.5 mL/min. Eluents: aqueous 0.1 % CF₃CO₂H/MeOH. Gradient: t = 0-5 min: 10 % MeOH; 5-30 min: 10→100 % MeOH; 30-34 min: 100 % MeOH; 34-35 min: 100→10 % MeOH; 35-40 min: 10 % MeOH. γ detection.




Figure S6. RP-HPLC analytical chromatograms of **Re1** (UV detection, 220 nm) and **Tc1** (γ -detection).