

**Select a page**

Clay Sampler [Samples of all our earths and clays](#) Kaolin [Our Kaolins and information](#) DE  
[Diatomaceous Earth information](#)  
[Diatomaceous earth Background](#) -- [Diatomaceous earth for human use](#) -- [Bentonite for Human Use](#) --  
[Bentonite Background](#)  
Must read links Excerpts from the Handbook of Clay Science [CLAYS AND HUMAN HEALTH](#)-- [CLAYS  
AND MINERALS AS DRUGS](#)  
Clay Home [Return to Homepage](#)

**Or continue reading****CLAYS AND CLAY MINERALS AS DRUGS**

M.T. DROY-LEFAIX<sup>a</sup> AND F. TATEO<sup>b</sup>

*<sup>a</sup>Beaufour-IPSEN, F-75016 Paris, France*

*<sup>b</sup>Istituto di Ricerca sulle Argille, CNR, I-85050 Tito Scalo (PZ), Italy*

(Please note that the terms bentonite and montmorillonite are synonyms.)

Montmorillonite is the main constituent of the volcanic ash weathering product, bentonite.)

Clay therapy is based on the ability of clays and clay minerals to adsorb and retain harmful and toxic substances. The beneficial effects of these materials to human health, notably in the treatment of gastrointestinal disorders, were recognized.

Indeed, the eating of clay ('geophagy') was practiced since antiquity in all parts of the world. Among the variety of clays and clay minerals that were used by primitive tribes are bentonite, kaolinite, montmorillonite, smectite (calcium and sodium bentonites), and 'pascalite'

(a calcium bentonite from Wyoming, USA) ([Eaton and Eaton, 1995](#)).

Examination of the diets of certain tribes in the high Andes of South America and central Africa, and those of Australian aborigines, showed that these people use clay to avoid getting stomach-ache, dysentery, and food infections. Indeed, the Quetchus Indians of South America used to dip their potatoes into an aqueous suspension of clay, immediately before eating, in order to prevent the build-up of acidity in the stomach. This dietetic procedure is still being followed by some tribes of American Indians. A similar practice was traditionally carried out on board ships where sailors

used clays not only to adsorb odours and moisture but also to treat dysentery, burns, boils, sore mouths, and other internal and external disorders.

Although recent research confirmed that clays and clay minerals possess general curative properties, it is the treatment of disorders that remains the focus of attention.

By adsorbing 'aggressors' (infectious factors) of the gastrointestinal mucosa barrier, these materials can serve as both prophylactic and therapeutic agents.

### **11.6.1. INTERACTIONS OF CLAY MINERALS WITH GASTROINTESTINAL MUCUS**

At the surface of the gut, a mucus gel adheres to the epithelial cells of the mucosa. This adherent mucus is dynamic, being continuously secreted by the caliciform cells and regularly eroded by environmental 'aggressors' present in the gut lumen.

The mucus gel is largely composed of glycoprotein polymers, lipids, and proteins, linked together by covalent bonds. As such, it acts as a physical barrier protecting the mucosa against penetration by extraneous molecules and mechanical injury.

By maintaining a pH gradient and competing with the epithelial surface for microorganisms,

the mucus gel also acts as a chemical barrier.

Thus, a weakening of the mucus gel barrier may be at the origin of disorders such as gastritis and colitis (Droy-Lefaix, 1987). Short-term treatment with clay minerals, such as smectite (calcium and sodium bentonites) (More' et al., 1987) and attapulgite (More' et al., 1992) increases the

thickness of the adherent mucus. This may be ascribed to interactions of mineral particles with mucus components (Leonard et al., 1994) by which the gastrointestinal glycoproteins are modified, and their polymerization is enhanced (Droy-Lefaix et al., 1986). Similarly, aluminium (hydr)oxides (e.g., boehmite) can reduce mucus degradation (Bouyssou et al., 1990). The beneficial effects of minerals are also associated

with improvements in the rheological properties of the mucus gel, such as spinability. This reflects the increased extent of polymerization, and the improvement in quality, of the adherent mucus (Droy-Lefaix et al., 1985). Changes in the physico-chemical properties of the mucus, induced by the action of clay minerals, were confirmed by electron paramagnetic resonance and fluorescence spectroscopy. The results indicate that clay mineral ingestion decreases mucus solubility. At the same time, the viscosity and hydrophobicity of the mucus increases, enhancing its adhesion to epithelial cells.

### **11.6.2. CLAY MINERALS, MUCOSAL BARRIER, AND GASTROINTESTINAL ‘AGGRESSORS’**

By acting directly on the mucus gel, clays and clay minerals exert a stabilizing effect on the mucosal barrier (Gwozdziński et al., 1997), providing protection against different ‘aggressors’ of the gastrointestinal mucosa.

Pepsin, a substance necessary for digestion, is a typical ‘aggressor’. Experiments with rats showed that if pepsin secretion at the surface of the gastric mucosa is strongly increased (due to pathological dysregulation), the adherent mucus layer is progressively disrupted. At the same time haemorrhagic mucosal lesions appear, and significant bleeding occurs in the lumen as well as localized ulceration in an otherwise intact epithelium. By binding to the mucus components, smectite (calcium and sodium bentonites) can completely

inhibit the damage induced by pepsin (Leonard et al., 1994).

Samson et al. (1995), for example, showed that patients with ulcerative colitis show a six-fold greater mean total faecal proteinase activity (expressed in mmol terminal NH<sub>2</sub>/min/g dry weight of faeces) than the control. smectite (calcium and sodium bentonites) totally inhibits

this enzyme activity. The effects of smectite (calcium and sodium bentonites) on mucus proteolysis are assessed using a

model of mucolytic activity, assayed by the release of degraded colonic mucin from the adherent mucus gel of freshly prepared pig colonic bags in vitro. Similarly, trypsin (2 mg/mL) releases three times more soluble mucin per bag than the control. smectite (calcium and sodium bentonites) (100 mg/mL) inhibits trypsin activity, causing the level of degraded

mucin to fall below the normal value. This is ascribed to the interaction of smectite (calcium and sodium bentonites)

with the adherent mucus layer, and the binding of trypsin to the mineral (Samson et al., 1995).

Clay minerals can also provide protection against attack by bile acids that cause gastrointestinal ulceration. In rats, oral administration of sodium glycodeoxycholate or sodium taurocholate induces severe erosion of the jejunal mucosa. After treatment with smectite (calcium and sodium bentonites) (which interacts closely with the mucus glycoproteins) the severity of

surface erosion is greatly diminished (Fioramonti et al., 1990), while the rheological properties of the adherent mucus gel are maintained within normal limits (Droy-Lefaix et al., 1985).

Because of their strong bioadhesive properties, clay minerals also afford protection of the colon against damage from reactive oxygen species. Oxygenated free radicals, released by infiltration of white cells into the colonic mucosa barrier, are very unstable. Their presence can induce severe erosion of the colonic mucosa, leading to mucolysis. By maintaining the solution viscosity of the colonic mucin, and inhibiting the hypersecretion of mucus, smectite (calcium and sodium bentonites) can prevent the onset of mucolysis

(Pearson et al., 1996; Knight et al., 1998).

In many digestive diseases, the intestinal barrier is weakened by the release of proinflammatory

cytokines, induced by abnormal activation of the epithelial cells and the underlying immune system. These cytokines include a tumour necrosis factor- $\alpha$  (TNF- $\alpha$ ) and an interferon- $\gamma$  factor (INF- $\gamma$ ). When intestinal cells (line HT 29-19 A) are incubated with TNF- $\alpha$  and INF- $\gamma$ , intestinal function (assessed in Ussing chambers by measuring ionic conductance, apicobasal fluxes of  $^{14}\text{C}$ -mannitol, and intact horseradish peroxidase) is altered, and the tight junction between cells is disrupted. In the presence of smectite (calcium and sodium bentonites) (100 mg/mL) the values of these parameters are similar to those of the control (Mahraoui et al., 1997).

The cytoprotective effects of clay minerals can also account for their ability to prevent damage of the gastrointestinal mucosa caused by such 'aggressors' as ethanol and anti-inflammatory drugs. Ethanol, directly administered into the stomach, gives rise to severe gastric ulcerations and macroscopic necrosis of the gastric mucosa. These deleterious effects are accompanied by a decrease in the gastric transmural potential difference which serves as a criterion of the functional integrity of the mucosa (Fioramonti et al., 1990). Erosion of the mucus layer leads to a significant alteration of rheological properties (Droy-Lefaix et al., 1992; Slitine-Bonet et al., 1994). smectite (calcium and sodium bentonites) treatment for two days can significantly counteract the harmful effect of ethanol, reducing the irritative index (Fioramonti et al., 1990).

Clay minerals also provide protection against the action of anti-inflammatory drugs. For example, the oral administration of aspirin (2 g) to pigs, and phenylbutazone (200 mg/kg) to rats, decreases the gastric potential difference, and induces severe ulceration due to mucus alteration (Fioramonti et al., 1990; Droy-Lefaix et al., 1992). The extent of lesion and mucus degradation is significantly reduced after treatment with smectite (calcium and sodium bentonites). In humans the symptoms of gastropathies, induced by

non-steroid anti-inflammatory drugs, can be successfully treated by ingestion of smectite (calcium and sodium bentonites) (Peignot et al., 1997).

### 11.6.3. ADSORPTIVE PROPERTIES OF CLAYS AND CLAY MINERALS

The adsorptive properties of clay minerals provide the basis for the therapeutic uses of clays.

#### A. Toxins

Clays can adsorb a variety of toxic substances, such as strychnine (Droy-Lefaix, 1986), mycotoxins (e.g., T2 toxin) (Fioramonti et al., 1987b), aflatoxin (Schell et al., 1993), enterotoxins (Brouillard and Rateau, 1989), and toxins produced by *Vibrio cholerae*, *Escherichia coli* (Fioramonti et al., 1987b), and *Yersinia pseudotuberculosis* (Carnoy et al., 2000).

By doing so, clays can provide active protection against disturbances during gastrointestinal transit. In mice, for example, gastric emptying and small intestinal transit are significantly accelerated after oral administration (1 mg/kg for 4 days) of T2 toxin. However, if the toxin is incubated with smectite (calcium and sodium bentonites) for 24 h beforehand, no increase in the rate of gastric emptying and small intestinal transit occurs (Fioramonti et al., 1987a).

In conscious dogs, intraduodenal administration of cholera toxin (200 mg) affects gastrointestinal transit, and disrupts the migrating motor complexes (MMCs) of the stomach and jejunum. According to the duration of treatment (at a dose of 100 mg/kg/day), smectite (calcium and sodium bentonites) can effectively counteract the effects of cholera toxin (Fioramonti et al., 1987b).

smectite (calcium and sodium bentonites) can also adsorb the enterotoxin of *Clostridium difficile* (Martirosian et al., 1998). In rats, this toxin causes intestinal permeability to increase through

hypersecretion of colonic water. Both these conditions can be alleviated by treatment with  $^{51}\text{Cr}$ -EDTA in the presence of smectite (calcium and sodium bentonites) (Fioramonti et al., 1994).

*E. coli* toxin is an infectious agent causing diarrhoea. Heat-stable toxin (ST) from *E. coli*, directly administered to New Zealand rabbits, induces a significant increase in intestinal permeability (as estimated by Evans Blue) and severe damage to ileal loops (as revealed by scanning electron microscopy). The presence of smectite (calcium and sodium bentonites) in the ileal loops has a protective effect (Pons et al., 1997).

Similar results are obtained with the enterotoxin of *Bacteroides fragilis* administered to HT/29 C1 cells (human colon adenocarcinoma cell line). Prior incubation of this toxin with smectite (calcium and sodium bentonites) suppresses its toxic effects (Martirosian et al., 1998).

### B. Pesticides

Because of their high adsorptive capacity, clay minerals can also protect the digestive mucosa against pesticide damage. Diquat, a widely used non-selective desiccant herbicide, induces erosion of intestinal mucosa and fluid hypersecretion. In rats that were given diquat, treatment with smectite (calcium and sodium bentonites) (500 mg/kg for 2 weeks) brings about a

normalization of mucus rheological properties and intestinal permeability, as

746 Chapter 11.6: Clays and Clay Minerals as Drugs

indicated by urine analysis using  $^{51}\text{Cr}$ -EDTA (Theodorou et al., 1995). Similarly, montmorillonite and bentonite are good adsorbents, and may be recommended for the treatment of pesticide poisoning (Meredith and Vale, 1987).

### C. Microorganisms

Clay minerals are efficient drugs for treating disorders of the gastrointestinal mucosa, induced by microorganisms. Kaolinite and montmorillonite are capable of

adsorbing viruses (Lipson and Stotzky, 1984). As such, these minerals can induce rapid recovery when administered to children suffering from gastroenteritis.

Similarly, the strong adsorptive power of smectite (calcium and sodium bentonites) lies behind its ability to aggregate

bacteria, such as strains of *E. coli* with the plasmid P, carrying a virulence factor in the form of an external protein CS 31A (Girardeau, 1987).

In the stomach, *Helicobacter pylori* is associated with gastritis and gastroduodenal ulcers. This bacterium is also one of the most important ethiopathogenic factors causing peptic ulcer. smectite (calcium and sodium bentonites), on HeLa cells infected by *H. pylori* isolated from

human biopsies, significantly reduces adhesion of the bacteria to the surface of epithelial cells (Bonneville et al., 1990). This is why smectite (calcium and sodium bentonites) is effective in treating

the symptoms of people with non-ulcer dyspepsia who are infected by *H. pylori* (De Korwin et al., 1993).

In the intestine, smectite (calcium and sodium bentonites) is effective against diarrhoea as shown by clinical data for

new-born calves with neonatal gastroenteritis. Faeces analysis reveals the presence of rotavirus in 41.3% of the animals as well as that of *E. coli* K99, coronavirus, and *Salmonella*. Recovery is observed in 72% of calves after 2.8 and 2.2 days of receiving 250 and 500mg/kg smectite (calcium and sodium bentonites), respectively, and after 4.2 days in calves which do not

receive smectite (calcium and sodium bentonites). After 4 days of treatment, the consistency of the faeces is significantly

better in calves receiving smectite (calcium and sodium bentonites) than in the control animals (Espinasse et al., 1987).

#### D. Gas

Clay minerals can serve as gas adsorbents in patients with symptoms of flatulence

and abdominal distension. Thus, smectite (calcium and sodium bentonites) can reduce the amount of hydrogen

emitted during colonic fermentation (Frexinos et al., 1986; Arbeille et al., 1991).

#### E. Alimentary Allergy

Food allergy is also responsible for disturbances in colonic transit, water absorption, and intestinal permeability. Guinea pigs that were sensitized by b-lactoglobulin from cow milk show colonic transit acceleration, a colonic hypersecretory response, a strong increase in intestinal permeability, and a decrease in faecal dry matter. These effects are not observed in animals that were treated with smectite (calcium and sodium bentonites). Clays can inhibit

anaphylaxia probably by controlling the release of mediators at the origin of the degranulation of the mast cells (Theodorou et al., 1994).

#### 11.6.4. CLAY MINERALS AND CLINICAL APPLICATIONS

Being good adsorbents and mucostabilizers, clay minerals are efficacious against several aggressive agents causing severe intestinal disorders. Acute gastroenteritis is a major cause of morbidity and mortality among children worldwide (Madkour et al., 1993). By adsorbing viruses, bacteria, and other digestive irritants, clay minerals can shorten the course of acute diarrhoea, and reduce the occurrence of prolonged diarrhoea.

Furthermore, these minerals do not interfere with the electrolyte balance, and are well tolerated by patients (Buttron, 1987; DuPont et al., 1990; Bauer and Hirschbrunn, 1992; Charritat et al., 1992; Dupont et al., 1992; Vivatvakin et al., 1992; Lexomboon et al., 1994; Karas, 1996; Milocco et al., 1999; Guarino et al., 2001; Narkeviciute et al., 2002).

Clay minerals also provide protection against diarrhoeas induced by antibiotics treatments (Benhamou et al., 1995), alleviate chronic diarrhoeas induced by chemotherapy

and radiation (Hornbrink et al., 1995; Ippolite, 1998; Santantonio et al.,

2000), enteral nutrition (Perrotin et al., 1990), and HIV infection (Phanuphak et al., 1992; Mastroianni et al., 1998).

Clay minerals are promising drugs in the treatment of irritable bowel syndrome (IBS), a rather frequent disease in adults with a complex pathogenic mechanism. By enhancing the thickness of the mucus barrier, both colon movement function and faeces consistency are restored, and the symptoms of IBS are alleviated (Opriu et al., 1996; Secondulfo et al., 2002). In parallel, clay minerals have a positive effect on flatulence and abdominal distension (Lukas and Lukas, 2000).

#### 11.6.5. CONCLUSIONS

Clay minerals protect and are efficient against several ‘aggressors’ that cause major disorders of the gut. These beneficial effects of clay minerals (on the gastrointestinal mucosa) are associated with two mechanisms of action: (1) adsorption of the ‘aggressors’

or their toxic secretions and (2) modification of the thickness and rheological properties of the adherent mucus, reinforcing the natural defenses of the gastrointestinal mucosa.

**Links to more information about our clays and earths**

**Clay Sampler [Samples of all our earths and clays](#)**

**Kaolin [Our Kaolins and kaolin information from the internet](#)**

**Diatomaceous Earth [Diatomaceous Earth information](#)**

**[Diatomaceous earth Background](#)**

**[Diatomaceous earth for human use](#)**

**[Bentonite for Human Use](#)**

**[Bentonite Background](#)**

**Clay Home [Return to Homepage](#)**

**These are must read links**

**Excerpts from the Handbook of Clay Science**

**[CLAYS AND HUMAN HEALTH](#)**

**[CLAYS AND MINERALS AS DRUGS](#)**