

# Heterochromia in Designed Metallopeptides: Geometry-Selective Binding of Cd<sup>II</sup> in a De Novo Peptide\*\*

Olga Iranzo, Chris Cabello, and Vincent L. Pecoraro\*

One of the most important chemical concepts is defining how one molecule recognizes and controls the properties of another molecule or ion. Nowhere is this issue more significant than in the field of biomolecular recognition. Metalloproteins efficiently control the geometry and coordination number of metal ions as well as the types of ligands bound to them. Incorporation of metals in the correct binding site is essential for proper biological activity of the protein, regardless of whether the function is catalytic, structural, or regulatory. Despite the importance of metalloproteins, the major factors determining metal-ion selectivity and specificity are not yet well understood. Recent studies on metalloregulatory proteins and carbonic anhydrase suggest that coordination number and geometry of the metal ion are key elements.<sup>[1,2]</sup> This control becomes even more significant when proteins can bind two metal ions with different coordination environments. This is the case for the enzyme 5'-aminolevulinic acid dehydratase, which contains two Zn<sup>II</sup> centers, one bound in a five-coordinate geometry and the other bound in a pseudotetrahedral geometry. Interestingly, lead toxicity is mainly due to the replacement by a Pb<sup>II</sup> ion of the Zn<sup>II</sup> ion bound to the tetrahedral site.<sup>[3,4]</sup> Thus, a deep knowledge of the factors determining this delicate ion recognition is crucial to understanding the principles that govern metalloprotein structure and function.

Using the de novo designed TRI peptide family (TRI = Ac-G-(LKALEEK)<sub>4</sub>-G-NH<sub>2</sub>), we have been investigating the influence of protein environment on metal-ion specificity and how subtle changes in the amino acid sequence can fully control the coordination number of the metal ion. By exploiting our previous observations that we could influence

the metal binding affinity in the TRI coiled coils by placing cysteine in either position **a** or **d** of the heptad repeat unit, we designed a TRI derivative with two cysteine binding sites (TRI L9CL19C, Table 1) that shows sequential and selective binding of Cd<sup>II</sup> ions, first to the **a** site and then to the **d** site.<sup>[5]</sup>

**Table 1:** Derivatives of the TRI peptide.

Peptide	Sequence <sup>[a]</sup>
TRI L9CL19C	Ac-G LKALEEK <b>CK</b> ALEEK LK <b>ACE</b> EK LKALEEK G-NH <sub>2</sub>
TRI L12AL16C	Ac-G LKALEEK LK <b>AA</b> EK <b>CK</b> ALEEK LKALEEK G-NH <sub>2</sub>
TRI L16Pen	Ac-G LKALEEK LKALEEK <b>XK</b> ALEEK LKALEEK G-NH <sub>2</sub>
TRI L12AL16CL23Pen	Ac-G LKALEEK LK <b>AA</b> EK <b>CK</b> ALEEK <b>XK</b> ALEEK G-NH <sub>2</sub>
TRI L16PenL23CL26A	Ac-G LKALEEK LKALEEK <b>XK</b> ALEEK <b>CK</b> AAEAK G-NH <sub>2</sub>
Grand L16PenL26AL30C	Ac-G LKALEEK LKALEEK <b>XK</b> ALEEK LK <b>AA</b> EK <b>CK</b> ALEEK G-NH <sub>2</sub>
Grand L26AL30C	Ac-G LKALEEK LKALEEK LKALEEK LK <b>AA</b> EK <b>CK</b> ALEEK G-NH <sub>2</sub>
Grand L16Pen	Ac-G LKALEEK LKALEEK <b>XK</b> ALEEK LKALEEK LKALEEK G-NH <sub>2</sub>

[a] X = Penicillamine. Residues in bold indicate modifications.

This result highlights how site-specific binding of Cd<sup>II</sup> ions can be achieved in a de novo designed peptide containing two binding sites that are defined by the same first-coordination-sphere ligands, three cysteine residues in both cases, solely by changing the conformation and environment of the metal-binding cavity. Later, using the **a** binding sites, we designed two peptides, TRI L16Pen and TRI L12AL16C (Table 1), that are able to bind Cd<sup>II</sup> ions exclusively in a trigonal-planar structure and a pseudotetrahedral geometry, respectively. This precise control of the coordination number and geometry of the Cd<sup>II</sup> centers was achieved by controlling the steric constraints in the metal-binding pocket.<sup>[6]</sup> We know that these complexes, [Cd(TRI L16Pen)<sub>3</sub>] and [Cd(TRI L12AL16C)<sub>3</sub>], have different physical properties, as shown by their UV and CD spectra at pH 8.5 ([Cd(TRI L16Pen)<sub>3</sub>] UV band:  $\lambda_{227}$  (28 900 m<sup>-1</sup> cm<sup>-1</sup>), CD band:  $\lambda_{227}$  (5000 deg dmol<sup>-1</sup> cm<sup>2</sup>) and  $\lambda_{257}$  (-3500 deg dmol<sup>-1</sup> cm<sup>2</sup>); [Cd(TRI L12AL16C)<sub>3</sub>] UV band:  $\lambda_{230}$  (21 800 m<sup>-1</sup> cm<sup>-1</sup>), CD band:  $\lambda_{237}$  (5500 deg dmol<sup>-1</sup> cm<sup>2</sup>) and  $\lambda_{260}$  (-590 deg dmol<sup>-1</sup> cm<sup>2</sup>)) and also by their different pH profiles for binding Cd<sup>II</sup> centers, as explained below.

While being able to impose a specific coordination number on a metal was a signal achievement, a more challenging construct would be a single polypeptide that utilizes subtle sequence perturbations to bind selectively the same metal into distinct coordination structures. Specifically, our next goal was to design a single short peptide capable of simultaneously binding two Cd<sup>II</sup> ions, one with a trigonal-planar geometry and the second one with a pseudotetrahedral geometry, within the same three-stranded coiled coil. While

[\*] Dr. O. Iranzo, C. Cabello, Prof. V. L. Pecoraro  
Department of Chemistry  
University of Michigan  
Ann Arbor, Michigan 48109-1055 (USA)  
Fax: (+1) 734-936-7628  
E-mail: vlpec@umich.edu  
Prof. V. L. Pecoraro  
Biophysics Research Division  
University of Michigan  
Ann Arbor, Michigan 48109-1055 (USA)

[\*\*] V.L.P. thanks the National Institute of Health for support of this research (R01 ES0 12236) and O.I. thanks the Margaret and Herman Sokol Foundation for a Postdoctoral Award.


other investigations have shown that de novo peptides can bind two metals in separate mononuclear sites<sup>[5,7]</sup> or in binuclear centers,<sup>[8–10]</sup> they have not addressed the question of selective binding of the same type of metal ion in different centers having distinct geometries. Two important issues to be addressed with our studies are whether the presence of one metal center influences the properties of a second center and whether different properties can be observed for the same metal in distinct sites. We designate heterochromic peptides as those capable of binding two equivalents of the same metal ion in a similar way while enforcing different physical properties in those metals. This is equivalent to an individual having two eyes, but of different color. These studies are of fundamental importance to understand the factors that control metal-ion specificity, an important objective of de novo designed metalloproteins.

Our initial strategy was to introduce the amino acid modifications that led to the trigonal planar {CdS<sub>3</sub>} unit and to the pseudotetrahedral {CdS<sub>3</sub>O} unit into the TRI peptide sequence. The first modification involves the replacement of the cysteine residue with penicillamine, and the second one calls for the substitution of the leucine residue above the cysteine layer for alanine.<sup>[6]</sup> The <sup>113</sup>Cd NMR spectrum of a solution containing our first construct, TRI L12AL16CL23Pen (Table 1), and two equivalents of Cd<sup>II</sup> at pH 9.6 shows two resonances with <sup>113</sup>Cd chemical shifts of  $\delta = 593$  ppm and 574 ppm. Based on our previous results,<sup>[6,11]</sup> these <sup>113</sup>Cd chemical shifts are consistent with a peptide that binds two Cd<sup>II</sup> ions, both as primarily the {CdS<sub>3</sub>O} species (Table 2). Thus, while we were successful obtaining the four-

coordinate site, we did not succeed in designing the three-coordinate center. We reasoned that if position 23 was too near the edge of the coiled coil and the coordination of water to the Cd<sup>II</sup> center were promoted by the fraying of the  $\alpha$ -helices, then moving the three-coordinate site towards the center of the coiled coil might avoid this situation and yield a binding site with a higher percentage of the three-coordinate species. Hence, we switched the positions of cysteine and penicillamine to obtain the new construct TRI L16PenL23-CL26A (Table 1). At pH 9.6, the <sup>113</sup>Cd NMR spectrum of a solution containing this peptide and two equivalents of Cd<sup>II</sup> shows two resonances with  $\delta = 676$  ppm and 609 ppm. These


chemical shifts are consistent with a peptide that binds two Cd<sup>II</sup> centers, one present to 90–95% as the {CdS<sub>3</sub>} species ( $\delta = 676$  ppm) and the other to approximately 80% as the {CdS<sub>3</sub>O} species ( $\delta = 609$  ppm; Table 2). These results indicate that to define the three-coordinate site, it is important to control not only the first coordination sphere around the Cd<sup>II</sup> center but also its location inside the coiled coil.

Our next goal was to improve the design of the four-coordinate site, keeping the two binding sites strictly independent. To achieve this aim, we moved to the Grand peptides, which contain an extra heptad repeat unit, thus allowing the introduction of the cysteine residue seven amino acids further down the sequence (L30C). It was felt that this new construct, Grand L16PenL26AL30C (Table 1), should tolerate the introduction of an alanine residue above the cysteine without disrupting the three-coordinate site. The corresponding <sup>113</sup>Cd NMR spectrum at pH 9.8 shows two resonances, one at  $\delta = 687$  ppm and the other at  $\delta = 588$  ppm (Figure 1). These results are consistent with a peptide that binds two Cd<sup>II</sup> ions, one with a pure trigonal planar {CdS<sub>3</sub>} coordination geometry ( $\delta = 687$  ppm) and the other with a pure pseudotetrahedral {CdS<sub>3</sub>O} geometry ( $\delta = 588$  ppm; Table 2).



**Figure 1.** <sup>113</sup>Cd NMR spectra of a solution containing 2.6 mM trimer Grand L16PenL26AL30C and 5.2 mM <sup>113</sup>Cd(NO<sub>3</sub>)<sub>2</sub> at different pH values.

Having succeeded in making a single, short peptide capable of binding Cd<sup>II</sup> ions in a trigonal S<sub>3</sub> environment in one position and in a pseudotetrahedral S<sub>3</sub>O environment in a second position, we next investigated the control of metal-ion binding and the impact on the physical properties of the centers. The pH dependence of the formation of the complex [Cd<sub>2</sub>(Grand L16PenL26AL30C)<sub>3</sub>]<sup>2-</sup> was monitored at 235 nm, and the pH titration curve is shown in Figure 2a. Interestingly, this UV/Vis pH titration reveals a very different pH profile for Cd<sup>II</sup> ions binding to each of the sites in Grand L16PenL26AL30C. Two well-defined regions are observed, and their shapes are consistent with the simultaneous release of two protons, as was observed previously for the TRI peptides and related monosubstituted peptides.<sup>[11,12]</sup> The best fit to the experimental data was obtained by analyzing each region independently and using the reported model, which is shown in Equation (1).<sup>[12]</sup>




**Table 2:** Percentages of the four-coordinate Cd<sup>II</sup> species {CdS<sub>3</sub>O} for different peptides of the TRI family.

Peptide	Four-coordinate Cd <sup>II</sup> {CdS <sub>3</sub> O} <sup>[a]</sup> [%]	
	L#AL#C or L#CL#A site	L#Pen site
TRI L12AL16C	100	–
TRI L16Pen	–	0
TRI L12AL16CL23Pen	100	90
TRI L16PenL23CL26A	80	5–10
Grand L16PenL26AL30C	100	0

[a] Values determined based on correlation between the percentages of three- and four-coordinate Cd<sup>II</sup> species from <sup>111m</sup>Cd PAC spectroscopy (PAC = perturbed angular correlation of  $\gamma$  rays) and the corresponding <sup>113</sup>Cd NMR chemical shifts obtained for different Cd<sup>II</sup> complexes of the TRI peptide family.<sup>[6,11]</sup>

coordinate site, we did not succeed in designing the three-coordinate center. We reasoned that if position 23 was too near the edge of the coiled coil and the coordination of water to the Cd<sup>II</sup> center were promoted by the fraying of the  $\alpha$ -helices, then moving the three-coordinate site towards the center of the coiled coil might avoid this situation and yield a binding site with a higher percentage of the three-coordinate species. Hence, we switched the positions of cysteine and penicillamine to obtain the new construct TRI L16PenL23-CL26A (Table 1). At pH 9.6, the <sup>113</sup>Cd NMR spectrum of a solution containing this peptide and two equivalents of Cd<sup>II</sup> shows two resonances with  $\delta = 676$  ppm and 609 ppm. These



**Figure 2.** UV/Vis pH titrations: a) pH dependence of 2 equiv (●) and 1 equiv (■) of Cd<sup>II</sup> binding to Grand L16PenL26AL30C (20  $\mu\text{M}$  trimer); b) pH dependence of 2 equiv of Cd<sup>II</sup> binding to Grand L16PenL26AL30C (20  $\mu\text{M}$  trimer; ●), of 1 equiv of Cd<sup>II</sup> binding to Grand L26AL30C (20  $\mu\text{M}$  trimer; ▲), and of 1 equiv of Cd<sup>II</sup> binding to Grand L16Pen (20  $\mu\text{M}$  trimer; ■).

The apparent  $\text{p}K_{\text{a}2}$  values determined for each region are reported in Table 3. Repetition of the experiment with only one equivalent of Cd<sup>II</sup> in solution indicates selective binding of Cd<sup>II</sup> ions to the lower  $\text{p}K_{\text{a}2}$  binding site (Figure 2a); however, UV spectral data alone can not establish which site corresponds to the {CdS<sub>3</sub>} or {CdS<sub>3</sub>O} geometries.

To assign binding sites to these different  $\text{p}K_{\text{a}2}$  values, we synthesized two Grand peptides, each containing only one Cd<sup>II</sup> binding site: Grand L26AL30C (four-coordinate site) and Grand L16Pen (three-coordinate site). The pH-dependent curves of Cd<sup>II</sup> binding to these Grand peptides are shown in Figure 2b. These pH profiles are also consistent with the

**Table 3:** Apparent  $\text{p}K_{\text{a}2}$  values for TRI and Grand peptides.

Peptide	Apparent $\text{p}K_{\text{a}2}$ <sup>[a]</sup>
TRI L12AL16C	12.2 $\pm$ 0.2
TRI L16Pen	15.3 $\pm$ 0.2
Grand L26AL30C	9.9 $\pm$ 0.2
Grand L16Pen	15.7 $\pm$ 0.2
Grand L16PenL26AL30C	9.6 $\pm$ 0.2 (L12AL30C site) 16.1 $\pm$ 0.2 (L16Pen site)


[a] The best fit to the experimental data is consistent with the model given in Equation (1), which corresponds to the simultaneous release of two protons.  $A = (c_M \epsilon) / (10^{(\text{p}K_{\text{a}2} - 2\text{pH})} + 1)$ , where  $A$  is the observed absorbance,  $c_M$  is the total metal concentration, and  $\epsilon$  is the extinction coefficient of the complex.<sup>[11,12]</sup>

simultaneous release of two protons, and the fit of the experimental data using the model mentioned above [Eq. (1)] give the apparent  $\text{p}K_{\text{a}2}$  values reported in Table 3. These titrations provide two useful pieces of information. First, the  $\text{p}K_{\text{a}2}$  values suggest that the low- $\text{p}K_{\text{a}2}$  species in Grand L16PenL26AL30C corresponds to the four-coordinate site ( $\text{p}K_{\text{a}2} = 9.6$  vs.  $\text{p}K_{\text{a}2} = 9.9$  for Grand L26AL30C) while the high  $\text{p}K_{\text{a}2}$  ( $\text{p}K_{\text{a}2} = 16.1$  vs.  $\text{p}K_{\text{a}2} = 15.7$  for Grand L16Pen) is associated with the three-coordinate Cd<sup>II</sup> center. Second, the close similarities between the  $\text{p}K_{\text{a}2}$ 's of Cd<sup>II</sup> ions binding to the monosubstituted peptides with those determined for the disubstituted peptide indicate that the two binding sites in Grand L16PenL26AL30C behave independently of each other.

To confirm our assignments, we carried out <sup>113</sup>Cd NMR spectroscopic pH titrations (Figure 1). At pH 6.6 only a single resonance is observed with a <sup>113</sup>Cd chemical shift of  $\delta = 588$  ppm, indicating binding of Cd<sup>II</sup> ions to the four-coordinate site. When the pH value is raised to 9.8, a second resonance appears at  $\delta = 687$  ppm, the hallmark signal for {CdS<sub>3</sub>}. These results corroborate the UV/Vis pH titrations and clearly show that the binding site capable of sequestering Cd<sup>II</sup> ions at low pH values corresponds to the site with {CdS<sub>3</sub>O} coordination. The apparent  $\text{p}K_{\text{a}2}$  value of 9.6 obtained for the Cd<sup>II</sup> ion binding to this site is the lowest observed to date for our TRI peptide family.

Because we know from the pH titrations that the two metal sites in Grand L16PenL26AL30C behave independently of one another, there must be another explanation for the lowering of the  $\text{p}K_{\text{a}2}$  value of the {CdS<sub>3</sub>O} site. The most reasonable candidate is the difference in affinity of the Grand versus TRI peptide frameworks. We determined the pH profile of Cd<sup>II</sup> ions binding to TRI L12AL26C and TRI L16Pen. Consistent with the previous data, two protons are released in each system; the apparent  $\text{p}K_{\text{a}2}$  values are reported in Table 3. All these data indicate that while the pH dependence of Cd<sup>II</sup> ions binding to the three-coordinate site {CdS<sub>3</sub>} is not affected to a large extent in moving from Grand to TRI peptides, the four-coordinate site is perturbed. This effect is most likely a reflection of different self-association affinities and pH-dependent aggregation state preferences of the peptides themselves. The Grand peptides show a higher self-association affinity than TRI peptides,<sup>[13]</sup> which could result in the formation of three-stranded coiled coils at pH values lower than 6.0. This observation suggests that the formation and stabilization of the four-coordinate {CdS<sub>3</sub>O} unit may be achieved at more acidic pH values than those reported herein by increasing the self-association energy of the three-stranded coiled coil relative to the two-stranded coiled coil.

Figure 3 shows the <sup>113</sup>Cd NMR spectra corresponding to the titration of successive aliquots of 0.5 equivalents of <sup>113</sup>Cd(NO<sub>3</sub>)<sub>2</sub> into a solution containing 2.6 mM trimer Grand L16PenL26AL30C at pH 9.6. A single peak at  $\delta = 588$  ppm is observed up to one equivalent of Cd<sup>II</sup>. A second resonance at  $\delta = 687$  ppm appears on addition of the second equivalent of Cd<sup>II</sup>. We conclude that under conditions in which Cd<sup>II</sup> ions will bind exclusively as either {CdS<sub>3</sub>O} (pseudotetrahedral) or {CdS<sub>3</sub>} (trigonal planar; i.e. pH > 9.0), these data show



**Figure 3.**  $^{113}\text{Cd}$  NMR spectra of solutions containing 2.6 mM trimer Grand L16PenL26AL30C and different amounts of  $^{113}\text{Cd}(\text{NO}_3)_2$  at pH 9.6.

unambiguously that this metal ion binds selectively to the four-coordinate site. Thus, regardless of the pH value, the four-coordinate site is the first filled by  $\text{Cd}^{\text{II}}$ . This result most likely reflects the preference of  $\text{Cd}^{\text{II}}$  for this coordination environment. These data, together with the pH titration experiments explained above, point out how  $\text{Cd}^{\text{II}}$  ions bound to these two binding sites possess different physical properties and how they behave independently of one another.

In conclusion, we are able to design a single, short peptide (Grand L16PenL26AL30C) capable of binding two  $\text{Cd}^{\text{II}}$  ions within the same three-stranded coiled coil, one with a pseudotetrahedral and the other with a trigonal planar coordination geometry. More interestingly, we have shown how  $\text{Cd}^{\text{II}}$  ions bound to the two different sites possess different physical properties. Binding of  $\text{Cd}^{\text{II}}$  ions to the four-coordinate site occurs at much lower pH value ( $\text{p}K_{\text{a}2} = 9.6$ ) than binding to the three-coordinate site ( $\text{p}K_{\text{a}2} = 16.1$ ). Moreover,  $\text{Cd}^{\text{II}}$  binds selectively to the four-coordinate site under all pH conditions. Thus, the heterochromic peptide Grand L16PenL26AL30C shows site-selective metal-ion recognition and controls the physical properties of  $\text{Cd}^{\text{II}}$  centers by controlling their coordination geometry. This advance portends a new generation of heterochromic metallo-biomolecules that may be used as selective bifunctional switches.

### Experimental Section

**Peptide synthesis and purification:** The different peptides of the TRI family (see Table 1 for sequences) were synthesized on an Applied Biosystems 433 A peptide synthesizer using standard protocols<sup>[14]</sup> and purified and characterized as described previously.<sup>[15]</sup>

**UV/Vis spectroscopy:** Fresh solutions of the purified peptides were prepared for each experiment using doubly distilled water. The peptide concentration of the solutions was determined by quantization of the cysteine thiol groups using 4,4'-dipyridyl disulphide.<sup>[16]</sup> The pH titrations were performed as previously described.<sup>[12]</sup> A concentration of 60  $\mu\text{M}$  monomer peptide and the corresponding amount of  $\text{CdCl}_2$  were used in all experiments. In all cases, reverse titrations were carried out to verify the reversibility of the process. The pH-dependent absorption spectra were fit using the same models and procedure as in our previous studies.<sup>[11,12]</sup>

**$^{113}\text{Cd}$  NMR spectroscopy:** All spectra were collected at room temperature on a Varian Inova 500 spectrometer (110.92 MHz for  $^{113}\text{Cd}$ ) equipped with a 5-mm broadband probe.  $^{113}\text{Cd}$  NMR spectra were externally referenced to a 0.1M  $\text{Cd}(\text{ClO}_4)_2$  solution in  $\text{D}_2\text{O}$ . A spectral width of 847 ppm (93 897 Hz) was sampled using a 5.0- $\mu\text{s}$  90° pulse and 0.05-s acquisition time with no delay between scans. Samples were prepared under a flow of argon by dissolving 30–35 mg of the lyophilized and degassed peptides in 450–500  $\mu\text{L}$  15%  $\text{D}_2\text{O}$  solution. The peptide concentrations were determined using 4,4'-dipyridyl disulphide.<sup>[16]</sup> The final samples were prepared by the addition of the appropriate amount of 250 mM  $^{113}\text{Cd}(\text{NO}_3)_2$  solution (prepared from 95% isotopically enriched  $^{113}\text{CdO}$  obtained from Oak Ridge National Laboratory), and the pH value was adjusted with KOH or HCl solution. An argon atmosphere was maintained when possible, but the samples came in contact with  $\text{O}_2$  while the pH value was adjusted. The actual final concentrations for each experiment are indicated in the text and the figure captions. The data were analyzed using the software MestRe-C.<sup>[17]</sup> All free induction decays (FIDs) were zero-filled to double the original points and were processed by application of 50-Hz line broadening prior to Fourier transformation.

Received: April 18, 2007

Published online: June 20, 2007

**Keywords:** cadmium · coordination modes · de novo peptides · metalloproteins · protein design

- [1] K. A. McCall, C. A. Fierke, *Biochemistry* **2004**, *43*, 3979, and references therein.
- [2] M. A. Pennella, A. I. Arunkumar, D. P. Giedroc, *J. Mol. Biol.* **2006**, *356*, 1124, and references therein.
- [3] P. T. Erskine, N. Senior, S. Awan, R. Lambert, G. Lewis, I. J. Tickle, M. Sarwar, P. Spencer, P. Thomas, M. J. Warren, P. M. Shoolingin-Jordan, S. P. Wood, J. B. Cooper, *Nat. Struct. Biol.* **1997**, *4*, 1025.
- [4] E. K. Jaffe, J. Martins, J. Li, J. Kervinen, R. L. J. Dunbrack, *J. Biol. Chem.* **2001**, *276*, 1531.
- [5] M. Matzapetakis, V. L. Pecoraro, *J. Am. Chem. Soc.* **2005**, *127*, 18229.
- [6] K.-H. Lee, C. Cabello, L. Hemmingsen, E. N. G. Marsh, V. L. Pecoraro, *Angew. Chem.* **2006**, *118*, 2930; *Angew. Chem. Int. Ed.* **2006**, *45*, 2864.
- [7] T. Tanaka, T. Mizuno, S. Fukui, H. Hiroaki, J.-I. Oku, K. Kanaori, K. Tajima, M. Shirakawa, *J. Am. Chem. Soc.* **2004**, *126*, 14023.
- [8] A. Lombardi, C. M. Summa, S. Geremia, L. Randaccio, V. Pavone, W. F. DeGrado, *Proc. Natl. Acad. Sci. USA* **2000**, *97*, 6298.
- [9] C. M. Summa, M. M. Rosenblatt, J.-K. Hong, J. D. Lear, W. F. DeGrado, *J. Mol. Biol.* **2002**, *321*, 923.
- [10] K. Spiegel, W. F. De Grado, M. L. Klein, *Proteins Struct. Funct. Bioinf.* **2006**, *65*, 317.
- [11] O. Iranzo, D. Ghosh, V. L. Pecoraro, *Inorg. Chem.* **2006**, *45*, 9959.
- [12] M. Matzapetakis, D. Ghosh, T.-C. Weng, J. E. Penner-Hahn, V. L. Pecoraro, *J. Biol. Inorg. Chem.* **2006**, *11*, 876.
- [13] D. Ghosh, K.-H. Lee, B. Demeler, V. L. Pecoraro, *Biochemistry* **2005**, *44*, 10732.
- [14] W. C. Chan, P. D. White, *Fmoc Solid Phase Peptide Synthesis: A Practical Approach*, Oxford University Press, New York, **2000**.
- [15] B. T. Farrer, N. P. Harris, K. E. Balchus, V. L. Pecoraro, *Biochemistry* **2001**, *40*, 14696.
- [16] M. Mantle, G. Stewart, G. Zayas, M. King, *Biochem. J.* **1990**, *266*, 597.
- [17] C. Cobas, J. Cruces, F. J. Sardina, MestRe-C version 2.3, Universidad de Santiago de Compostela, Spain, **2000**.